

A Web of Worlds presents  
The Ultimate Cheat Sheet for Astrophysics Students

Lachlan Marnoch  
[www.webofworlds.net](http://www.webofworlds.net)

v 1.0.1  
July 31, 2018

# Contents

<b>1</b>	<b>Physics</b>	<b>5</b>
1.1	Motion	5
1.1.1	Velocity	5
1.1.2	Acceleration	5
1.1.3	Newton's Laws	5
1.1.4	Momentum	5
1.1.5	Centripetal Force	5
1.1.6	Kinetic Energy	5
1.1.7	Projectile Motion	6
1.1.8	Rotation	6
1.1.9	Euler-Lagrange and the Hamiltonian	7
1.2	Oscillations	7
1.2.1	Springs	7
1.3	Materials	8
1.3.1	Density	8
1.4	Energy	8
1.4.1	Work	8
1.5	Forces	8
1.5.1	Buoyancy (Archimedes' Principle)	8
1.5.2	Friction	8
1.6	Waves	8
1.6.1	Wavelength	8
1.6.2	Angular Frequency	8
1.7	Newtonian Gravity	8
1.7.1	Force of Gravity	8
1.7.2	Gravitational Potential (potential energy per unit mass)	8
1.7.3	Gravitational field	8
1.7.4	Gravitational Potential Energy	8
1.7.5	Kepler's Third Law	9
1.8	Electromagnetism	9
1.8.1	Notation	9
1.8.2	Maxwell's Equations	9
1.8.3	Lorentz Force	9
1.8.4	Electric Field	9
1.8.5	Dipole moment	10
1.8.6	Electric potential	10
1.8.7	Electric potential difference	10
1.8.8	Electric potential energy	10
1.8.9	Charge densities	10
1.8.10	Current densities	10
1.8.11	Circuits	11
1.8.12	Capacitors	11
1.8.13	Magnetic fields	12
1.8.14	Inductors	12
1.8.15	Materials	12
1.9	Special Relativity	13

1.9.1	Interval	13
1.9.2	Four-vectors	14
1.9.3	Frames of Reference	15
1.9.4	Proper Velocity	15
1.10	General Relativity	15
1.10.1	Metrics	15
1.10.2	Rindler coordinates	16
1.10.3	Einstein notation	16
1.10.4	Christoffel symbols	16
1.10.5	Covariant derivatives	17
1.10.6	Riemann curvature tensor	17
1.10.7	Ricci curvature tensor	17
1.10.8	Einstein's equations	17
1.11	Thermodynamics	17
1.11.1	Ideal Gases	17
1.11.2	Microstates	17
1.11.3	Entropy	18
1.11.4	Black bodies	18
1.12	Quantum Mechanics	18
1.12.1	The Uncertainty Principle	18
1.12.2	Bras and Kets	18
1.12.3	Rules for an Inner Product	18
1.12.4	The Born Rule	18
1.12.5	Expectation	18
1.12.6	Variance	19
1.12.7	Standard Deviation	19
1.12.8	Trace	19
1.12.9	Partial Trace	19
1.12.10	The Schrödinger Equation	19
1.12.11	Heisenberg equation of motion	19
1.12.12	Operators	19
<b>2</b>	<b>Astrophysics &amp; Astronomy</b>	<b>22</b>
2.1	Astrometry	22
2.1.1	Redshift	22
2.1.2	Apparent magnitude	22
2.1.3	Absolute magnitude	22
2.1.4	Relative magnitudes	22
2.1.5	Flux-magnitude relationship	22
2.1.6	Color	22
2.1.7	Metallicity	22
2.2	Stars	22
2.2.1	Stellar Structure Equations	22
2.2.2	Timescales	23
2.2.3	Gravitational potential energy	23
2.2.4	Eddington Limit (hydrostatic equilibrium)	23
2.2.5	Mass-Luminosity Relationship	23
2.3	Galaxies	23
2.3.1	Hubble Elliptical Galaxy Classification	23
2.3.2	Sérsic Profile	24
2.3.3	Density of stars in the Milky Way Galaxy	24
2.4	Black Holes	24
2.4.1	Schwarzschild Radius	24
2.5	Instrumentation	24
2.5.1	Lensmaker's equation	24
2.5.2	Focal ratio / Focal number	24
2.5.3	Field of view	24
2.5.4	Resolution Limits	24

2.5.5	Nyquist sampling	24
2.5.6	Plate scale	24
2.5.7	Fitting error	25
2.5.8	Adaptive optics error	25
2.5.9	Signal-to-noise ratio	25
2.5.10	Atmospheric Extinction	25
2.5.11	Rocket science	25
<b>3</b>	<b>Mathematics</b>	<b>26</b>
3.1	Notation	26
3.2	Algebra	26
3.2.1	Factorisation	26
3.2.2	Absolute Value	26
3.2.3	Quadratics	26
3.2.4	Logarithms	27
3.2.5	Vectors	27
3.2.6	Factorials	27
3.2.7	Inner product definition	28
3.2.8	Complex Numbers	28
3.2.9	Power Series	28
3.2.10	Matrix Operations	28
3.2.11	Matrix Types	29
3.2.12	Change of Basis Unitary	32
3.2.13	Commutator	32
3.2.14	Anticommutator	32
3.2.15	Cauchy-Schwarz Inequality	32
3.3	Geometry	32
3.3.1	Pythagorean theorem	32
3.3.2	Properties of shapes	32
3.3.3	Properties of solids	32
3.3.4	Circular formulae	33
3.3.5	Useful Functions	33
3.3.6	Coordinates	34
3.3.7	Hyperbolic Functions	34
3.4	Trigonometry	35
3.4.1	Identities	35
3.5	Calculus	39
3.5.1	Limits	39
3.5.2	Properties	39
3.5.3	Differentiation	39
3.5.4	Partial Differentiation	41
3.5.5	The Differential	41
3.5.6	Line Element	41
3.5.7	Integration	41
3.5.8	Vector Calculus	43
3.5.9	Dirac Delta Function	44
3.5.10	Approximations	44
<b>4</b>	<b>Statistics</b>	<b>45</b>
4.1	Variance	45
4.2	Standard Deviation	45
4.2.1	Population Standard Deviation	45
4.2.2	Sample Standard Deviation	45
4.3	Residual Sum of Squares	45
4.4	Mean Squared Error	45
4.5	Residual Standard Error	45
4.6	Correlation	45

<b>A</b>	<b>Values</b>	<b>46</b>
A.1	Physics	46
A.1.1	Physical Constants	46
A.1.2	Useful Quantities	47
A.2	Astronomy	47
A.2.1	Useful Quantities	47
A.3	Mathematics	48
<b>B</b>	<b>Units of Measurement</b>	<b>49</b>
B.1	Natural Units	49
B.2	SI System	49
B.2.1	Base Units	49
B.2.2	Derived Units	50
B.3	CGS (centimetres-grams-seconds)	51
B.4	Astronomy units	52
B.4.1	Astronomical system	52
B.4.2	Equatorial Coordinate System	52
B.5	United States customary units (aka Imperial Units)	53
B.5.1	Length	53
B.6	Degrees of Angle	53
B.7	Miscellaneous Units	53
B.7.1	Pressure	53
B.8	Prefixes	53
<b>C</b>	<b>Mathematical Stuff</b>	<b>55</b>
C.1	Trigonometric Values	55
C.1.1	Pythagorean Triples	55
<b>D</b>	<b>Boring stuff</b>	<b>56</b>
D.1	Version History	56
D.2	Licensing	56
D.3	Contact	56
D.4	Credits	56

# Chapter 1

## Physics

### 1.1 Motion

#### 1.1.1 Velocity

- $\vec{v} = \frac{\Delta \vec{x}}{\Delta t} = \frac{d\vec{x}}{dt} = \dot{\vec{x}}$

#### 1.1.2 Acceleration

- $\vec{a} = \frac{\Delta \vec{v}}{\Delta t} = \frac{d\vec{v}}{dt} = \frac{d^2 \vec{x}}{dt^2} = \ddot{\vec{x}}$

#### 1.1.3 Newton's Laws

##### Newton's First Law

- When viewed in an inertial reference frame, an object either remains at rest or continues to move at a constant velocity, unless acted upon by a net force.

##### Newton's Second Law

- $\vec{F}_{net} = m\vec{a} = \frac{d\vec{p}}{dt}$

##### Newton's Third Law

- $\vec{F}_A = -\vec{F}_B$
- When one body exerts a force on a second body, the second body simultaneously exerts a force equal in magnitude and opposite in direction on the first body.

#### 1.1.4 Momentum

- $\vec{p} = \gamma m \vec{v} \approx m \vec{v}$
- $\Delta \vec{p} = \vec{F} \Delta t$
- $\vec{F} = \frac{\Delta \vec{p}}{\Delta t} = \frac{d\vec{p}}{dt}$

#### 1.1.5 Centripetal Force

- $F_c = \frac{mv^2}{r}$

#### 1.1.6 Kinetic Energy

- $K = \frac{1}{2}mv^2$

### 1.1.7 Projectile Motion

- $v_y^2 = u_y^2 + 2a_y\Delta y$
- $x = u_x t$
- $\Delta y = u_y\Delta t + \frac{1}{2}a_y\Delta t^2 = u_y t + \frac{1}{2}\frac{F_y}{m}\Delta t^2$

### 1.1.8 Rotation

#### Angular Velocity

- $\omega = \frac{d\theta}{dt} = \dot{\theta}$
- $\omega = \frac{v}{r}$
- $\vec{v} = \vec{r} \times \vec{\omega}$

#### Angular Acceleration

- $\alpha = \frac{d\omega}{dt} = \frac{d^2\theta}{dt^2} = \dot{\omega} = \ddot{\theta}$

#### Moment of Inertia

##### *Point Mass*

- $I = mr^2$

##### *Several Point Masses*

- $I = \sum mr^2$

##### *Continuous mass*

- $I = \int r^2 dm$

##### *Parallel axis theorem*

- $I = I_{com} + md^2$

##### *Thin disc rotating about centre*

- $I = \frac{MR^2}{2}$

##### *Thin hoop rotating about centre*

- $I = MR^2$

##### *Thin rod rotating about centre*

- $I = \frac{ML^2}{12}$

##### *Thin rod rotating about end*

- $I = \frac{ML^2}{3}$

#### Rotational Kinetic Energy

- $K_{rot} = \frac{1}{2}I\omega^2$

#### Total Kinetic Energy

- $K_{tot} = K_{trans} + K_{rot} = \frac{1}{2}(mr_{com}^2 + I_{com})\omega^2$

## Angular Momentum

- $\vec{L} = I\vec{\omega} = \vec{r} \times \vec{p}$

## Torque

- $\vec{\tau} = I\vec{\alpha} = \frac{dL}{dt} = \vec{r} \times \vec{F}$

## 1.1.9 Euler-Lagrange and the Hamiltonian

### Lagrangian

- $\ell = T - V = \sum_{lm} a(q)\dot{q}_l\dot{q}_m$

- $= K(\dot{q}) - U(q)$

*Generalised coordinates & momenta*

- $p_k \equiv \frac{\partial L}{\partial \dot{q}_k}$

### Euler-Lagrange Equation

- $\frac{d}{dt} \frac{\partial \ell}{\partial \dot{x}} - \frac{\partial \ell}{\partial x} = 0$

### Action

- $S[x(t)] = \int_{t_A}^{t_B} \ell(\dot{x}(t), x(t)) dt$

### Hamiltonian

- $\mathcal{H} = \sum_i p_i \dot{q} - L$

- $\dot{P} = -\frac{\partial H}{\partial Q}$

- $\dot{Q} = \frac{\partial H}{\partial P}$

- $\dot{P} = -\omega^2 Q$

- $\dot{Q} = P$

## 1.2 Oscillations

### 1.2.1 Springs

#### Force of a Spring

- $\vec{F} = -k_s \vec{x}$

#### Potential Energy of a Spring

- $U_s = \frac{1}{2} k_s x^2$

#### Angular Frequency of a Spring

- $\omega = \sqrt{\frac{k_s}{m}}$



## 1.3 Materials

### 1.3.1 Density

- $\rho = \frac{m}{V} = \frac{dm}{dV}$

## 1.4 Energy

### 1.4.1 Work

- $W = \int_a^b \vec{F} \cdot d\vec{l} \approx \vec{F} \cdot \vec{s}$

## 1.5 Forces

### 1.5.1 Buoyancy (Archimedes' Principle)

- $F_{buoy} = m_{displaced}g = \rho_d V_d g$

### 1.5.2 Friction

- $F_K \approx \mu_K F_{\perp}$
- $F_S \leq \mu_S F_{\perp}$

## 1.6 Waves

- $a \sin(\omega t - kx + \phi)$
- $k = \frac{2\pi}{\lambda}$

### 1.6.1 Wavelength

- $v = f\lambda$

### 1.6.2 Angular Frequency

- $\omega = \frac{2\pi}{T} = 2\pi f$

## 1.7 Newtonian Gravity

### 1.7.1 Force of Gravity

- $\vec{F}_G = \frac{GmM}{r^2} \hat{r} = -m\vec{\nabla}\Phi(\vec{r}) \approx -mg\hat{y} = m\vec{g}$

### 1.7.2 Gravitational Potential (potential energy per unit mass)

- $\Phi(\vec{r}) = -\sum_i \frac{GM(\vec{r}_i)}{|\vec{r} - \vec{r}_i|} = -\int \frac{G\mu(\vec{r}')}{|\vec{r} - \vec{r}'|} d^3r'$

### 1.7.3 Gravitational field

- $\vec{g}(\vec{r}) = \frac{GM}{r^2} = -\nabla\Phi(\vec{r})$

### 1.7.4 Gravitational Potential Energy

- $U_G = -\frac{GmM}{r} \approx mgh$

### 1.7.5 Kepler's Third Law

- $\frac{T^2}{r^3} = \frac{4\pi^2}{G(m+M)} = \text{constant}$

## 1.8 Electromagnetism

### 1.8.1 Notation

- $\vec{z} = \vec{r} - \vec{r}'$

### 1.8.2 Maxwell's Equations

	Integral form	Differential form
Gauss's Law	$\oiint_S \vec{E} \cdot d\vec{a} = \frac{1}{\epsilon_0} \iiint_V \rho \, dV$ $= \frac{\sum Q_{enc}}{\epsilon_0}$	$\vec{\nabla} \cdot \vec{E} = \frac{\rho}{\epsilon_0}$
Gauss's Law for Magnetism	$\oiint_S \vec{B} \cdot d\vec{a} = 0$	$\vec{\nabla} \cdot \vec{B} = 0$
Maxwell-Faraday equation	$\oint_b \vec{E} \cdot d\vec{l} = -\frac{d}{dt} \iint_S \vec{B} \cdot d\vec{a}$	$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$
Ampère's circuital law	$\oint_b \vec{B} \cdot d\vec{l} = \mu_0 \iint_S \vec{J} \cdot d\vec{a} + \mu_0 \epsilon_0 \frac{d}{dt} \iint_S \vec{E} \cdot d\vec{a}$ $= \mu_0 (I_{enc} + \epsilon_0 \frac{d}{dt} \int_S \vec{E} \cdot d\vec{a})$	$\vec{\nabla} \times \vec{B} = \mu_0 (\vec{J} + \epsilon_0 \frac{\partial \vec{E}}{\partial t})$

### 1.8.3 Lorentz Force

On a point charge

- $\vec{F} = q(\vec{E} + \vec{v} \times \vec{B})$

On a current

- $d\vec{F} = I \int d\vec{l} \times \vec{B}$
- $\vec{F} = \vec{I}L \times \vec{B}$

### 1.8.4 Electric Field

- $\vec{E} = \int_V \frac{\rho(\vec{r}')}{z^2} \hat{z} \, d\tau$

From a single point charge

- $\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{r}$

From a dipole

- $|\vec{E}_{axis}| \approx \frac{2p}{4\pi\epsilon_0 r^3}$
- $|\vec{E}_{\perp}| \approx \frac{p}{4\pi\epsilon_0 r^3}$

### 1.8.5 Dipole moment

- $\vec{p} = q\vec{d}$

### 1.8.6 Electric potential

- $V = \frac{1}{4\pi\epsilon_0} \frac{Q}{z}$
- $\nabla^2 V = \frac{-\rho}{\epsilon_0}$

In a single-point charge field

- $\Delta(\vec{r}) = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$

### 1.8.7 Electric potential difference

- $\Delta(\vec{r}) = -\int_{\vec{b}}^{\vec{a}} \vec{E} \cdot d\vec{l}$

In a single-point charge field

- $\Delta(\vec{r}) = \frac{1}{4\pi\epsilon_0} Q \left( \frac{1}{b} - \frac{1}{a} \right)$

### 1.8.8 Electric potential energy

- $U_E = q\Delta V = \frac{1}{4\pi\epsilon_0} \frac{qQ}{z}$

Energy stored in an electrostatic field distribution

- $U_E = \frac{1}{2} = \epsilon_0 E^2 \times volume$

### 1.8.9 Charge densities

Surface

- $\sigma = \frac{dq}{da} = \frac{Q}{A}$

Line

- $\lambda = \frac{dq}{dl} = \frac{Q}{L}$

### 1.8.10 Current densities

Volume

- $\vec{J} = \frac{d\vec{I}}{d\vec{a}_{\perp}} = \frac{I}{A_{\perp}} = \sigma(\vec{E} + \vec{v} \times B) = |q|nu(\vec{E} + \vec{v} \times B)$
- $\vec{\nabla} \cdot \vec{J} = 0$

## Surface

- $$\vec{K} = \frac{d\vec{I}}{d\vec{l}_\perp} = \frac{I}{l} = \sigma\vec{v}$$

### 1.8.11 Circuits

#### Electron drift velocity

- $$\vec{v} = u\vec{E}_{net}$$

#### Current per unit charge

- $$i = nA_{cs}\vec{v} = nA_{cs}uE_{net}$$

#### Current

- $$I = ei = enA_{cs}uE_{net} = \frac{dq}{dt}$$

#### Electrical Power

- $$P = IV = I^2R$$

#### Voltage (Electric potential difference)

- $$V = \Delta V = IR = -\varepsilon$$

#### Electromotive Force (EMF) from a Non-Coulomb force

- $$\epsilon = \frac{F_{NC}d}{e}$$

#### Resistance

- $$R = \frac{L\rho}{A} = \frac{L}{\sigma A}$$
- $$R_{series} = R_1 + R_2 + \dots + R_n$$
- $$\frac{1}{R_{parallel}} = \frac{1}{R_1} + \frac{1}{R_2} + \dots + \frac{1}{R_n}$$

### 1.8.12 Capacitors

#### Capacitance

- $$C = \frac{Q}{V} = \frac{\varepsilon A}{d} = \frac{k\varepsilon_0 A}{d}$$

#### Energy stored in a capacitor

- $$W = \frac{CV^2}{2}$$

#### Electric field in a capacitor

- $$E = \frac{Q}{\varepsilon_0 A}$$

#### Potential difference across a capacitor

- $$\Delta V = -\frac{dQ}{A\varepsilon_0}$$

### 1.8.13 Magnetic fields

- $\vec{B}(\vec{z}) = \frac{\mu_0}{4\pi} \int \frac{\vec{I} \times \hat{z}}{z^2} dl$
- $d\vec{B} = \frac{\mu_0}{4\pi} \frac{q\vec{v} \times \hat{z}}{z^2} = \frac{\mu_0}{4\pi} \frac{I d\vec{l} \times \hat{z}}{z^2}$

Magnetic field due to a wire

- $\vec{B} = \frac{\mu_0}{4\pi} \frac{2I}{r} \hat{\phi}$

Magnetic vector potential

- $\vec{A}(\vec{z}) = \frac{\mu_0}{4\pi} \int \frac{\vec{J}(\vec{r}')}{z} d\tau$
- $\vec{\nabla} \times \vec{A} = \vec{B}$
- $\vec{\nabla} \times (\vec{\nabla} \times \vec{A}) = -\mu_0 \vec{J}$
- $\vec{\nabla} \cdot \vec{A} = 0$

### 1.8.14 Inductors

- $\varepsilon = -LI$

Energy stored in an inductor

- $W = \frac{LI^2}{2}$

### 1.8.15 Materials

Macroscopic Maxwell's Equations (Materials)

	Integral form	Differential form
Gauss's Laws	$\oiint_S \vec{P} \cdot d\vec{a} = -\sum Q_B$ $\oiint_S \vec{D} \cdot d\vec{a} = \sum Q_f$	$\vec{\nabla} \cdot \vec{P} = -\rho_B$ $\vec{\nabla} \cdot \vec{D} = \rho_f$
Gauss's Law for Magnetism	$\oiint_S \vec{B} \cdot d\vec{a} = 0$	$\vec{\nabla} \cdot \vec{B} = 0$
Maxwell-Faraday equation	$\oint_b \vec{E} \cdot d\vec{l} = -\frac{d}{dt} \iint_S \vec{B} \cdot d\vec{a}$	$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$
Ampère's circuital law	$\oint_b \vec{H} \cdot d\vec{l} = I_{f,enc} + \frac{\partial}{\partial t} \iint_S \vec{D} \cdot d\vec{a}$	$\vec{\nabla} \times \vec{H} = \vec{J}_f + \frac{\partial \vec{D}}{\partial t}$

Dielectric constant

- $k = \frac{\varepsilon}{\varepsilon_0} = \varepsilon_r$
- $\varepsilon = k\varepsilon_0 = \varepsilon_r \varepsilon_0$

Susceptibility

- $\chi_e = 1 - \varepsilon_r$

### Polarisability

- $\vec{P} = \epsilon_0 \chi_e \vec{E} = n\vec{p}$

### Bound Charge

Surface

- $\sigma_B = \vec{p} \cdot \hat{n}$

Volume

- $\rho_B = -\vec{\nabla} \cdot \vec{P}$

Total

- $Q_B = \sigma_B + \rho_B = \vec{p} \cdot \hat{n} - \vec{\nabla} \cdot \vec{P}$

### Electric displacement

- $\vec{D} = \epsilon \vec{E} = k\epsilon_0 \vec{E} = \epsilon_0 \vec{E} + \vec{P}$

### Magnetic field

- $\vec{H} = \frac{\vec{B}}{\mu_0} - \vec{M}$

### Magnetic dipole

- $\vec{m} = I\vec{a}$

### Bound current

- $\vec{J}_B = \vec{\nabla} \times \vec{M}$
- $\vec{K}_B = \vec{M} \times \hat{n}$

## 1.9 Special Relativity

### 1.9.1 Interval

- $\Delta s^2 = -c^2 \Delta t^2 + \Delta x^2 + \Delta y^2 + \Delta z^2$
- $ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2$
- $\Delta s^2 < 0$  is a timelike interval. Events separated by this interval can be causally related.
- $\Delta s^2 = 0$  is a lightlike interval. Events separated by this interval can be causally related, but only by a lightspeed signal.
- $\Delta s^2 > 0$  is a spacelike interval. Events separated by this interval CANNOT be causally related.

### Gamma Factor

- $\gamma = \frac{1}{\sqrt{1 - (\frac{v}{c})^2}}$
- $\gamma = \frac{dt}{d\tau}$

### Mass-energy

- $E_{rest} = mc^2$
- $E = \gamma mc^2 = \frac{1}{\sqrt{1 - (\frac{v}{c})^2}} mc^2$

### Relativistic kinetic energy

- $K = \gamma mc^2 - mc^2$

### Length contraction

- $l_v = \frac{l_0}{\gamma} = l_0 \sqrt{1 - (\frac{v}{c})^2}$

### Time dilation

- $t_v = \gamma t_0 = \frac{t_0}{\sqrt{1 - (\frac{v}{c})^2}}$

### Mass dilation

- $m_v = \gamma m_0 = \frac{m_0}{\sqrt{1 - (\frac{v}{c})^2}}$

### Relative Velocity

- $u'_x = \frac{\Delta x'}{\Delta t} = \frac{u_x - v_x}{1 - \frac{v_x u_x}{c^2}}$

### Relativistic Momentum

- $\vec{p} = \gamma \vec{v} = \frac{m\vec{v}}{\sqrt{1 - (v/c)^2}}$

## 1.9.2 Four-vectors

### Four-space

- $\mathbf{s} = \mathbf{x} = \begin{bmatrix} ct \\ x \\ y \\ z \end{bmatrix}$

### Four-velocity

- $\mathbf{u} = \frac{d\mathbf{s}}{d\tau} = \gamma \begin{bmatrix} c \\ v_x \\ v_y \\ v_z \end{bmatrix}$

### Four-momentum

- $\mathbf{p} = \begin{bmatrix} E/c \\ p_x \\ p_y \\ p_z \end{bmatrix} = \gamma m \begin{bmatrix} c \\ v_x \\ v_y \\ v_z \end{bmatrix} = m\mathbf{u}$

### 1.9.3 Frames of Reference

#### Condition for an inertial frame

- $\frac{d^2x}{dt^2} = \frac{d^2y}{dt^2} = \frac{d^2z}{dt^2} = 0$

#### Galilean Transformations

- $x' = x + vt$
- $y' = y$
- $z' = z$
- All assuming  $x$  is along the axis of motion and  $x = x'$  when  $t = 0$ .

#### Lorentz Boosts

- $t' = \gamma(t - \frac{vx}{c^2})$
- $x' = \gamma(x - vt)$
- $y' = y$
- $z' = z$
- ( $x$  is along the axis of motion)

- $$\begin{bmatrix} ct' \\ x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} \gamma & -v\gamma & 0 & 0 \\ -v\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} ct \\ x \\ y \\ z \end{bmatrix}$$

#### General Lorentz transformation

- $$\begin{bmatrix} b'^0 \\ b'^1 \\ b'^2 \\ b'^3 \end{bmatrix} = \begin{bmatrix} \gamma & -v\gamma & 0 & 0 \\ -v\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} b^0 \\ b^1 \\ b^2 \\ b^3 \end{bmatrix}$$
- Motion along the  $x$ -axis.

#### Proper Time

- $\tau = \int_{t_A}^{t_B} \frac{1}{\gamma} dt = \int_{t_A}^{t_B} \sqrt{1 - \frac{v^2(t)}{c^2}} dt$

### 1.9.4 Proper Velocity

- $\mathbf{u} = \frac{d\mathbf{s}}{d\tau}$

## 1.10 General Relativity

### 1.10.1 Metrics

#### Minkowski

- $$\eta = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$
- $ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2$



## Schwarzschild

- $$g = \begin{bmatrix} -(1 - \frac{2GM}{c^2 r}) & 0 & 0 & 0 \\ 0 & (1 - \frac{2GM}{c^2 r})^{-1} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{bmatrix}$$
- $$ds^2 = -(1 - \frac{2GM}{c^2 r})c^2 dt^2 + (1 - \frac{2GM}{c^2 r})^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$$

### 1.10.2 Rindler coordinates

#### Line element

- $$ds^2 = -(1 + \frac{gx'}{c^2})^2 (c dt')^2 + dx'^2$$

### 1.10.3 Einstein notation

- Contravariant:  $e^\alpha$
- Covariant:  $e_\alpha$
- $$t_{\alpha\beta} = g_{\beta\gamma} t_\alpha^\gamma$$
- $$t_\alpha^\beta = g^{\beta\gamma} t_{\alpha\gamma}$$
- $$t'^\alpha{}_\beta = \frac{\partial x'^\alpha}{\partial x^\gamma} \frac{\partial x^\delta}{\partial x'^\beta} t'^\gamma{}_\delta$$
- $$t'^\alpha{}_\beta = \frac{\partial x^\gamma}{\partial x'^\alpha} \frac{\partial x'^\delta}{\partial x^\beta} t_{\gamma\delta}$$

#### Metrics

- $$ds^2 = g_{\alpha\beta} dx^\alpha dx^\beta$$
- $$g^{\alpha\beta} = \frac{1}{g_{\alpha\beta}}$$
- $$\delta_\beta^\alpha = \begin{cases} 1 & \alpha = \beta \\ 0 & \alpha \neq \beta \end{cases}$$
- $$\delta_\gamma^\alpha a^\gamma = a^\alpha$$
- $$g^{\alpha\gamma} g_{\gamma\beta} = \delta_\beta^\alpha$$

#### Four-vector product

- $$\mathbf{a} \cdot \mathbf{b} = g_{\alpha\beta} a^\alpha b^\beta = a_\beta b^\beta$$

### 1.10.4 Christoffel symbols

- $$\Gamma^\alpha{}_{\beta\gamma} = \frac{1}{2} g^{\alpha\delta} \left( \frac{\partial g_{\delta\beta}}{\partial x^\gamma} + \frac{\partial g_{\delta\gamma}}{\partial x^\beta} - \frac{\partial g_{\beta\gamma}}{\partial x^\delta} \right)$$
- $$\Gamma_{\alpha\beta\gamma} = \frac{1}{2} \left( \frac{\partial g_{\delta\beta}}{\partial x^\gamma} + \frac{\partial g_{\delta\gamma}}{\partial x^\beta} - \frac{\partial g_{\beta\gamma}}{\partial x^\delta} \right)$$
- $$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu{}_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} = 0$$

### 1.10.5 Covariant derivatives

- $\nabla_\gamma t^\alpha{}_\beta = \frac{\partial t^\alpha{}_\beta}{\partial x^\gamma} + \Gamma^\alpha{}_{\gamma\delta} t^\delta{}_\beta - \Gamma^\delta{}_{\gamma\beta} t^\alpha{}_\delta$
- $\nabla_\gamma t^{\alpha\beta} = \frac{\partial t^{\alpha\beta}}{\partial x^\gamma} + \Gamma^\alpha{}_{\gamma\delta} t^{\delta\beta} + \Gamma^\beta{}_{\gamma\delta} t^{\alpha\delta}$
- $\nabla_\gamma t_{\alpha\beta} = \frac{\partial t_{\alpha\beta}}{\partial x^\gamma} - \Gamma^\delta{}_{\gamma\alpha} t_{\delta\beta} - \Gamma^\delta{}_{\gamma\beta} t_{\alpha\delta}$
- $\nabla_\gamma t_\alpha{}^\beta = \frac{\partial t_\alpha{}^\beta}{\partial x^\gamma} - \Gamma^\delta{}_{\gamma\alpha} t_{\delta}{}^\beta + \Gamma^\beta{}_{\gamma\delta} t_\alpha{}^\delta$

### 1.10.6 Riemann curvature tensor

- $R^\alpha{}_{\beta\gamma\delta} = \frac{\partial \Gamma^\alpha{}_{\beta\delta}}{\partial x^\gamma} - \frac{\partial \Gamma^\alpha{}_{\beta\gamma}}{\partial x^\delta} + \Gamma^\alpha{}_{\gamma\epsilon} \Gamma^\epsilon{}_{\beta\delta} - \Gamma^\alpha{}_{\delta\epsilon} \Gamma^\epsilon{}_{\beta\gamma}$
- $R_{\alpha\beta\gamma\delta} = \frac{1}{2} \left( \frac{\partial^2 g_{\alpha\delta}}{\partial x^\beta \partial x^\gamma} - \frac{\partial^2 g_{\alpha\gamma}}{\partial x^\beta \partial x^\delta} - \frac{\partial^2 g_{\beta\delta}}{\partial x^\alpha \partial x^\gamma} \right) + \frac{\partial^2 g_{\beta\gamma}}{\partial x^\alpha \partial x^\delta}$
- $R_{\alpha\beta\gamma\delta} = -R_{\beta\alpha\gamma\delta}$
- $R_{\alpha\beta\gamma\delta} = -R_{\beta\alpha\delta\gamma}$
- $R_{\alpha\beta\gamma\delta} = R_{\delta\gamma\alpha\beta}$
- $R_{\alpha\beta\gamma\delta} + R_{\alpha\delta\beta\gamma} + R_{\alpha\gamma\delta\beta} = 0$

### 1.10.7 Ricci curvature tensor

- $R_{\alpha\beta} = R^\gamma{}_{\alpha\gamma\beta}$
- $R = R^\alpha{}_\alpha$

### 1.10.8 Einstein's equations

- $R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$

## 1.11 Thermodynamics

### 1.11.1 Ideal Gases

#### Ideal Gas Law

- $pV = Nk_B T$

#### Heat / Thermal Energy

- $Q = mc\Delta T$

#### Heat Capacity

- $C = \frac{dQ}{dT}$

#### Specific Heat Capacity

- $c = \frac{C}{m}$

### 1.11.2 Microstates

- $\Omega = \frac{(q + N - 1)}{q!(N - 1)}$

### 1.11.3 Entropy

- $S = k_B \ln \Omega$

### 1.11.4 Black bodies

Energy of a photon

- $E = hf$

Wien's Displacement Law

- $\lambda_{max} = \frac{b}{T} = (2.8977729 \times 10^{-3}) \frac{1}{T}$

Stefan-Boltzmann Law

- $I = \sigma T^4$

Spectrum

- $B_\lambda(T) = \frac{2hc^2}{\lambda^5} \frac{1}{\exp(\frac{hc}{\lambda k_B T} - 1)}$
- $B_\nu(T) = \frac{2h\nu}{c^2} \frac{1}{\exp(\frac{h\nu}{k_B T}) - 1}$

## 1.12 Quantum Mechanics

### 1.12.1 The Uncertainty Principle

- $\Delta x \Delta p \geq \frac{\hbar}{2}$
- $\Delta E \Delta t \geq \frac{\hbar}{2}$

### 1.12.2 Bras and Kets

- $|\psi\rangle = \langle\psi|^\dagger$

### 1.12.3 Rules for an Inner Product

- $\langle\psi|\phi\rangle \equiv (|\psi\rangle, |\phi\rangle)$
- Symmetric:  
 $\langle\psi|\phi\rangle = \langle\phi|\psi\rangle^*$
- Linear in second component
- Anti-linear in first component

### 1.12.4 The Born Rule

- $P = |\langle\psi|\psi\rangle|^2$

### 1.12.5 Expectation

- $\langle A \rangle = \int A |\Psi(x, t)|^2 dx$
- $\langle A \rangle = \langle\psi|A|\psi\rangle$

### 1.12.6 Variance

- $\text{var}(A) = \langle \psi | A^2 | \psi \rangle - \langle \psi | A | \psi \rangle^2$

### 1.12.7 Standard Deviation

- $\delta A = \sqrt{\text{var}(A)} = \sqrt{\langle \psi | A^2 | \psi \rangle - \langle \psi | A | \psi \rangle^2}$

### 1.12.8 Trace

- $\text{Tr}(A) = \sum_j \langle x_j | A | x_j \rangle$

### 1.12.9 Partial Trace

- $\text{Tr}_B(|a\rangle\langle a| \otimes |b\rangle\langle b|) \equiv |a\rangle\langle a| \text{Tr}(|b\rangle\langle b|)$
- $\text{Tr}(k_{AB}) = \text{Tr}_A(\text{Tr}_B(k_{AB})) = \text{Tr}_B(\text{Tr}_A(k_{AB}))$
- $\rho_B = \text{Tr}_A(\rho_{AB})$
- The partial trace is linear

### 1.12.10 The Schrödinger Equation

- $i\hbar \frac{\partial}{\partial t} \Psi(r, t) = \hat{H} \Psi(r, t)$
- $-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi(x, t)}{\partial x^2} + V(x) \Psi(x, t) = i\hbar \frac{\partial \Psi(x, t)}{\partial t}$
- $-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x)}{\partial x^2} + V(x) \psi(x, t) = E \psi(x)$
- $\hat{H} |\Psi(t)\rangle = i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle$

### 1.12.11 Heisenberg equation of motion

- $\frac{d}{dt} \hat{A}(t) = \frac{i}{\hbar} [\hat{H}, \hat{A}(t)]$

### 1.12.12 Operators

- $a_{jk} = \langle j | A | k \rangle$

#### Diagonalizable Operator

- $A = \sum_j \lambda_j |\lambda_j\rangle\langle\lambda_j|$

#### Normal Operator

- $A = \sum_j |\lambda_j\rangle\langle\lambda_j|$

#### Eigenstate Operators

- $(|\lambda_k\rangle\langle\lambda_k|)^n = |\lambda_k\rangle\langle\lambda_k|$

#### Identity

- $I = \sum_j |x_j\rangle\langle x_j|$

### Projector

- $P = |\psi\rangle\langle\psi|$

### Density operator

- $\rho \equiv \sum_j P_j |\psi_j\rangle\langle\psi_j|$
- Hermitian:  $\rho^\dagger = \rho$
- Normalised:  $\text{Tr}(\rho) = 1$
- Positive Semi-Definite:  $\langle\psi|\rho|\psi\rangle \geq 0, \forall |\psi\rangle \in \mathbf{H}$
- $\text{purity} = \text{Tr}(\rho^2)$ 
  - $\frac{1}{d} \leq \text{Tr}(\rho^2) \leq 1$
  - Pure:  $\text{Tr}(\rho^2) = 1$
  - Maximally mixed:  $\text{Tr}(\rho^2) = \frac{1}{d}$
- $\rho_A = \text{Tr}_B(\rho_{AB})$
- $\langle A \rangle = \text{Tr}(\rho A)$

### Pauli Operators

- $\sigma_x = X = |0\rangle\langle 1| + |1\rangle\langle 0| \doteq \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
- $\sigma_y = Y = i|1\rangle\langle 0| - i|0\rangle\langle 1| \doteq \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$
- $\sigma_z = Z = |0\rangle\langle 0| - |1\rangle\langle 1| \doteq \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$
- $I = |0\rangle\langle 0| + |1\rangle\langle 1| \doteq \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  (Sometimes included)
- $\text{Tr} X = \text{Tr}(Y) = \text{Tr}(Z) = 0$
- With respect to Hilbert-Schmidt Inner Product:  
 $\|X\| = \|Y\| = \|Z\| = \|I\| = \sqrt{2}$

#### *Properties*

- Unitary
- Hermitian
- $\lambda = \pm 1$

### Photon Annihilation and Creation Operators

- $\hat{a}|n\rangle = \sqrt{n}|n-1\rangle$
- $\hat{a}^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle$
- $\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$
- $\langle\alpha|\hat{a}^\dagger = \alpha^*\langle\alpha|$

### Atomic Energy Level Operators (for a two-level approximation)

- $\hat{\sigma}_+ = |e\rangle\langle g|$
- $\hat{\sigma}_- = |g\rangle\langle e|$
- $\hat{\sigma}_z = |e\rangle\langle e| - |g\rangle\langle g|$
- $\hat{\sigma}_+|g\rangle = |e\rangle$
- $\hat{\sigma}_-|e\rangle = |g\rangle$
- $\hat{\sigma}_+|e\rangle = 0$
- $\hat{\sigma}_-|g\rangle = 0$

## Chapter 2

# Astrophysics & Astronomy

### 2.1 Astrometry

#### 2.1.1 Redshift

- $\frac{\lambda_{obs} - \lambda_{emit}}{\lambda_{emit}} \approx \frac{v}{c}$
- $1 + z = \frac{\lambda_{obs}}{\lambda_{emit}}$

#### 2.1.2 Apparent magnitude

- $m - m_0 = -2.5 \log_{10}\left(\frac{F}{F_0}\right)$

#### 2.1.3 Absolute magnitude

- $M = m - 5 \log_{10}\left(\frac{d}{10}\right)$

#### 2.1.4 Relative magnitudes

- $\frac{I_a}{I_b} = 100^{\frac{(m_b - m_a)}{5}}$

#### 2.1.5 Flux-magnitude relationship

- $F = F_0 \times 10^{-0.4m}$

#### 2.1.6 Color

- $-2.5 \log\left(\frac{F_{f1}}{F_{f2}}\right)$

#### 2.1.7 Metallicity

- $Z = \log_{10}\left(\frac{(Fe/H)_*}{(Fe/H)_{\odot}}\right) = \log_{10}(Fe/H)_* - \log_{10}(Fe/H)_{\odot}$

### 2.2 Stars

#### 2.2.1 Stellar Structure Equations

Hydrostatic Equilibrium

- $\frac{dP}{dr} = \frac{-GM_r \rho(r)}{r^2}$

### Mass Conservation

- $\frac{M_r}{r} = 4\pi r^2 \rho$

### Energy Equation

- $\frac{dL_r}{dr} = 4\pi r^2 \rho \epsilon$

### Radiative Transport

- $\frac{dT}{dr}|_{rad} = \frac{3}{4ac} \frac{\bar{\kappa} \rho}{T^3} \frac{L_r}{4\pi r^2}$

## 2.2.2 Timescales

### Thermal / Kelvin-Helmholtz Timescale

- $\tau_{KH} = \frac{|U_*|}{L_*} = \frac{GM_*^2}{R_* L_*}$
- $\tau_{KH\odot} \approx 50 \text{ million years}$

### Dynamical Timescale

- $\tau_{dyn} \approx \sqrt{\frac{R^3}{GM}} \approx \sqrt{G\bar{\rho}}$

### Nuclear Timescale / Main Sequence Lifespan

- $\tau_N \approx \tau_{\odot} M^{-3} \approx 10^9 \left(\frac{M}{M_{\odot}}\right)^{-3}$

## 2.2.3 Gravitational potential energy

- $U_* \approx \frac{-GM^2}{R}$

## 2.2.4 Eddington Limit (hydrostatic equilibrium)

- $L_{edd} = \frac{4\pi GMm_p c}{\sigma T} \approx 3.2 \times 10^4 \left(\frac{M}{M_{\odot}}\right) [L_{\odot}]$
- $M_{edd} = 3.1 * 10^{-5} \left(\frac{L}{L_{\odot}}\right) [M_{\odot}]$

### Eddington Rate (mass loss)

- $\dot{M}_{edd} = \frac{L_{edd}}{\eta c^2} \approx 2.4 \times 10^{-8} \left(\frac{M}{M_{\odot}}\right) [M_{\odot}/yr]$

## 2.2.5 Mass-Luminosity Relationship

- $\frac{L}{L_{\odot}} \approx b \left(\frac{M}{M_{\odot}}\right)^a ; \quad a, b = \begin{cases} 2.3, 0.23 & M < 0.43M_{\odot} \\ 4, 1 & 0.43M_{\odot} < M < 2M_{\odot} \\ 3.5, 1.5 & 2M_{\odot} < M < 20M_{\odot} \\ 1, 32000 & M > 55M_{\odot} \end{cases}$

## 2.3 Galaxies

### 2.3.1 Hubble Elliptical Galaxy Classification

- $10 \times \left(\frac{a-b}{a}\right)$



### 2.3.2 Sérsic Profile

- $I(R) = I_0 \exp\{-b[(\frac{R}{R_e})^{\frac{1}{n}} - 1]\}$

### 2.3.3 Density of stars in the Milky Way Galaxy

- $\rho(R, z) = \rho_0 e^{-z/z_0} e^{-R/h}$

## 2.4 Black Holes

### 2.4.1 Schwarzschild Radius

- $r_S = \frac{2GM}{c^2}$

## 2.5 Instrumentation

### 2.5.1 Lensmaker's equation

- $$\frac{1}{f} = (n-1)\left[\frac{1}{R_1} - \frac{1}{R_2} + \frac{(n-1)d}{nR_1R_2}\right]$$
$$\approx (n-1)\left[\frac{1}{R_1} - \frac{1}{R_2}\right] \quad (\text{Thin lens approximation})$$

### 2.5.2 Focal ratio / Focal number

- $N = \frac{f}{D}$

### 2.5.3 Field of view

- $FOV = \frac{w_D}{D_T N} = \frac{w_D}{f_{sys}}$

### 2.5.4 Resolution Limits

Diffraction limit (Rayleigh criterion)

- $\varepsilon_d = 1.22 \frac{\lambda}{D}$

Seeing limit (Rayleigh criterion)

- $\varepsilon_s = 0.98 \frac{\lambda}{r_0}$

Total Resolution limit

- $\varepsilon \sqrt{\varepsilon_d^2 + \varepsilon_s^2}$

### 2.5.5 Nyquist sampling

- $\frac{2p}{f_{sys}} = \frac{\lambda}{D_T}$  (When diffraction limited)

- $N = \frac{2p}{\lambda}$

### 2.5.6 Plate scale

- $\frac{1}{f} [\text{rad}/m] = \frac{206265}{f} [\text{arcsec}/m]$

### 2.5.7 Fitting error

- $\sigma_{fit}^2 = a_f \left(\frac{d_{sub}}{r_0}\right)^{\frac{5}{3}} = 0.26 \left(\frac{d_{sub}}{r_0}\right)^{\frac{5}{3}}$

### 2.5.8 Adaptive optics error

- $\sigma_{total}^2 = 0.3 \left(\frac{d_{sub}}{r_0}\right)^{\frac{5}{3}} + \left(\frac{\theta}{\theta_0}\right)^{\frac{5}{3}} + 28.4 \left(\frac{\tau}{\tau_0}\right)^{\frac{5}{3}} + C_{WFS} \left(\frac{\lambda}{F\tau d_{sub}}\right)^2$

### 2.5.9 Signal-to-noise ratio

- $SNR = \frac{Ft}{\sqrt{Ft + (B_s n_p t) + (D n_p t) + (R^2 n_p)}}$

### 2.5.10 Atmospheric Extinction

- $m_\lambda = m_{\lambda,z} - a_\lambda(\sec z)$

### 2.5.11 Rocket science

Tsiolkovsky rocket equation

- $\Delta v = v_e \ln\left(\frac{m_0}{m_f}\right)$

# Chapter 3

## Mathematics

### 3.1 Notation

- $[f(x)]_b^a = f(a) - f(b)$

### 3.2 Algebra

#### 3.2.1 Factorisation

- $(a + b)^2 = a^2 + b^2 + 2ab$
- $(a - b)^2 = a^2 + b^2 - 2ab$
- $a^2 - b^2 = (a + b)(a - b)$
- $(a + b)(a + c) = a^2 + (b + c)a + bc$
- $(a + b)^3 = a^3 + 3ab^2 + 3a^2b + b^3$
- $(a - b)^3 = a^3 + 3ab^2 - 3a^2b - b^3$
- $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
- $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$
- $a^{2n} - b^{2n} = (a^n - b^n)(a^n + b^n)$

#### 3.2.2 Absolute Value

- $|ab| = |a||b|$
- $\left|\frac{a}{b}\right| = \frac{|a|}{|b|}$
- $|a + b| \leq |a| + |b|$

#### 3.2.3 Quadratics

##### Quadratic Formula

For  $ax^2 + bx + c = 0$ ,  $a \neq 0$ :

- $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
- $b^2 - 4ac > 0$  - two real unequal solutions.
- $b^2 - 4ac = 0$  - repeated real solution.
- $b^2 - 4ac < 0$  - two complex solutions.

### 3.2.4 Logarithms

- $y = \log_b(x); x = b^y$
- $\log_b(xy) = \log_b x + \log_b(y)$
- $\log_b\left(\frac{x}{y}\right) = \log_b x - \log_b(y)$
- $\log_b(x^p) = p \log_b x$
- $\log_b(b^x) = x$
- $\log_b(a) = \frac{\log_d(a)}{\log_d(b)}$
  
- $\log_b(\sqrt[p]{x}) = \frac{1}{p} \log_b x$
- $p \log_b x + q \log_b(y) = \log_b(x^p y^q)$
- $b^{\log_b x} = x$
- $\log_b(b) = 1$
- $\log_b(1) = 0$

### 3.2.5 Vectors

#### Dot Product

- $\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3 + \dots + a_n b_n = ab \cos \theta$

#### Cross Product

- $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2 b_3 - a_3 b_2) \hat{i} + (a_3 b_1 - a_1 b_3) \hat{j} + (a_1 b_2 - a_2 b_1) \hat{k} = ab \sin \theta \hat{n}$
- $\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$
- $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$
- $\vec{a} \times \vec{b} = 0$

### 3.2.6 Factorials

- $n! = n(n-1)(n-2)\dots(2)(1)$
- $(n+1)! = (n+1)n!$

#### Stirling's approximation

- $n! \approx n \ln(n) - n + O(\ln(n))$

#### The Factorial Integral

- $\int_0^{\infty} x^n e^{-x} dx$

### 3.2.7 Inner product definition

1. Linear in first variable:

$$(\alpha a + \beta b, c) = \alpha(a, c) + \beta(b, c)$$

2. Positive-definite:

$$(a, a) \geq 0, (a, a) = 0 \iff a = 0$$

3. Conjugate symmetrical:

$$(a, b) = (b, a)^*$$

$$(a, b) = (b, a), b, a \in \mathbf{R}$$

### 3.2.8 Complex Numbers

- $z = a + ib = \Re(z) + \Im(z)i$

#### Euler's Formula

- $e^{i\theta} = \cos \theta + i \sin \theta$
- $re^{i\theta} = |z|e^{i\theta} = r(\cos \theta + i \sin \theta)$

#### De Moivre's Formula

- $(\cos x + i \sin x)^n = \cos(nx) + i \sin(nx)$

#### Complex Modulus

- $r = |z| = |a + ib| = \sqrt{a^2 + b^2} = \sqrt{\Re^2(z) + \Im^2(z)}$

#### Complex Conjugate

- $\bar{z} = (a + ib) = a - ib$
- $(a + ib)(a - ib) = |a + ib|^2$

#### Complex Argument

- $\theta = \arg(z) = \arctan\left(\frac{a}{b}\right) = \arctan\left(\frac{\Im(z)}{\Re(z)}\right)$

### 3.2.9 Power Series

- $f(x) = \sum_{n=0}^{\infty} f_n x^n$

#### Notable Series

- $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \dots$
- $\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$
- $\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$

### 3.2.10 Matrix Operations

#### Determinant

- $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$

### Transpose

- $a_{jk}^T = a_{kj}$
- $(AB)^T = B^T A^T$
- Linear:  $(A + B)^T = A^T + B^T$
- $(rA)^T = rA^T$

### Hermitian Adjoint

- $A^\dagger \equiv (A^*)^T = (A^T)^*$
- $(AB)^\dagger = B^\dagger A^\dagger$
- $(A + B)^\dagger = A^\dagger + B^\dagger$

### Trace

- $\text{Tr}(A) = \sum_j^n a_{jj}$
- Cyclic:  $\text{Tr}(AB) = \text{Tr}(BA)$
- Linear:  $\text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B)$   
 $\text{Tr}(aB) = a\text{Tr}(B), a \in \mathbf{C}$
- $\text{Tr}(SAS^{-1}) = \text{Tr}(A)$

### Hilbert-Schmidt Inner Product

- $(A, B) \equiv \text{Tr}(A^\dagger B)$

### Rank-Nullity Theorem

- $\text{rk}(A) = \dim(\ker(A)) + \dim(\text{im}(A))$

## 3.2.11 Matrix Types

### Real Matrix

- $a_{jk} \in \mathbf{R}$

### Square Matrix

- $m = n$

### Symmetric Matrix

- $A = A^T$
- $a_{jk} = a_{kj}$
- Square

### Normal Matrix

- $A^\dagger A = AA^\dagger$
- Square
- Diagonalisable

### Diagonal Matrix

- $a_{jk} = 0, j \neq k$
- $\lambda_j = a_{jj}$
- Square
- $e^D = \begin{bmatrix} e^{d_{11}} & 0 & \dots & 0 \\ 0 & e^{d_{11}} & \dots & 0 \end{bmatrix}$

### Diagonalisable Matrix

- $A = PDP^{-1}$
- Square

### Identity Matrix

- $IA = A$
- $i_{jj} = 1$
- $i_{jk} = 0, j \neq k$
- Real
- Square
- Diagonal
- Symmetric
- Hermitian

### Hermitian Matrix

- $H = H^\dagger$
- $h_{jk} = h_{kj}^*$
- $h_{jj} \in \mathbf{R}$
- $\lambda \in \mathbf{R}$
- Square
- Normal
- All real, square matrices are Hermitian

### Anti-Hermitian Matrix

- $H = -H^\dagger$
- $H_{jk} = -H_{kj}^*$
- Square

### Orthogonal Matrix

- $A^T = A^{-1}$
- $AA^T = I$
- $(AA^T)_{jk} = \delta_{jk}$

### Positive Semidefinite

- $A \geq 0$
- $\hat{A}^\dagger = \hat{A}, \hat{A} \geq 0$
- $B = \hat{A}^\dagger \hat{A}$  is positive semidefinite for any linear operator  $\hat{A}$
- Positive semidefinite matrices are Hermitian

### Projector

- $\hat{P}^2 = \hat{P}$
- $\lambda = 1$  or  $0$
- $P_1 P_2 \mapsto \mathbf{H}_1 \cap \mathbf{H}_2$
- Projectors are Hermitian

### Real Matrix

- $A = A^*$
- $A_{jk} = A_{jk}^*$

### Imaginary Matrix

- $A = -A^*$
- $A_{jk} = -A_{jk}^*$

### Symmetric Matrix

- $A = A^T$
- $A_{jk} = A_{kj}$
- Square

### Antisymmetric Matrix

- $A = -A^T$
- $a_{jk} = -a_{kj}$
- $a_{jj} = 0$
- Square

### Unitary Matrix

- $U^\dagger U = U U^\dagger = I$
- $U^\dagger = U^{-1}$
- $(U^\dagger U)_{jk} = \delta_{jk}$
- Square
- Normal
- Hermitian



### 3.2.12 Change of Basis Unitary

- $(V)_b = [U^\dagger]_a (V)_a$
- $[U]_a = [(b_0)_a (b_1)_a \dots (b_n)_a]$

### 3.2.13 Commutator

- $[A, B] = AB - BA$
- $[A, A] = 0$
- $[A + B, C] = [A, C] + [B, C]$
- $[A, BC] = [A, B]C + B[A, C]$

### 3.2.14 Anticommutator

- $\{A, B\} = AB + BA$

### 3.2.15 Cauchy-Schwarz Inequality

- $|\langle \vec{u}, \vec{v} \rangle|^2 \leq \langle \vec{u}, \vec{u} \rangle \cdot \langle \vec{v}, \vec{v} \rangle$

## 3.3 Geometry

### 3.3.1 Pythagorean theorem

- $a^2 + b^2 = c^2$
- $a = \sqrt{b^2 + c^2}$

#### Higher dimensions

- $r = \sqrt{x^2 + y^2 + z^2}$
- $r = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \sqrt{\vec{r} \cdot \vec{r}}$

#### Distance between two points

In two dimensions

- $d_{12} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

In higher dimensions

- $d_{ab} = \sqrt{(b_1 - a_1)^2 + (b_2 - a_2)^2 + \dots + (b_n - a_n)^2}$

### 3.3.2 Properties of shapes

	Area	Circumference
Circle	$\pi R^2$	$2\pi R$
Square	$L^2$	$4L$

### 3.3.3 Properties of solids

	Surface Area	Volume
Sphere	$4\pi R^2$	$\frac{4}{3}\pi R^3$

### 3.3.4 Circular formulae

Arc length

- $l = R\theta$

Area of a sector

- $A = \frac{R^2\theta}{2}$

Area of a segment

- $A = \frac{R^2}{2}(\theta - \sin \theta)$

### 3.3.5 Useful Functions

Parabola

- $f(x) = a(x - h)^2 + k$
- Vertex at  $(h, k)$
- Up-concave if  $a > 0$ ; down-concave if  $a < 0$
  
- $f(x) = ax^2 + bx + c$
- Vertex at  $(-\frac{b}{2a}, f(-\frac{b}{2a}))$
- Up-concave if  $a > 0$ ; down-concave if  $a < 0$

Hyperbola

- $(\frac{x - h}{a})^2 - (\frac{y - k}{b})^2 = 1$
- Centre at  $(h, k)$
- Asymptotes through centre, slope  $\pm \frac{b}{a}$

Circle

- $(x - h)^2 + (y - k)^2 = R^2$
- Centre at  $(h, k)$

Ellipse

- $1 = (\frac{x - h}{a})^2 + (\frac{y - k}{b})^2$
- Centre at  $(h, k)$
- Vertices  $a$  units right/left from the centre and vertices  $b$  units up/down from the center.

Sphere

- $R^2 = (x - h)^2 + (y - k)^2 + (z - l)^2$
- Centre at  $(h, k, l)$ :

### Ball

- $R^2 < (x - h)^2 + (y - k)^2 + (z - l)^2$
- Centre at  $(h, k, l)$ :

### 3.3.6 Coordinates

#### Transformations to Cartesian coordinates

<b>Cartesian</b>	$x = x$	$y = y$	$z = z$	$dV = dx dy dz$
<b>Polar (2D)</b>	$x = r \cos \phi$	$y = r \sin \phi$	N/A	$dA = r dr d\phi$
<b>Cylindrical</b>	$x = r \cos \phi$	$y = r \sin \phi$	$z = z$	$dV = r dr d\theta dz$
<b>Spherical</b>	$x = r \sin \theta \cos \phi$	$y = r \sin \theta \sin \phi$	$z = r \cos \theta$	$dV = r^2 \sin \theta dr d\theta d\phi$

#### Transformations from Cartesian coordinates

<b>Cartesian</b>	$x = x$	$y = y$	$z = z$
<b>Polar (2D)</b>	$r = \sqrt{x^2 + y^2}$	$\phi' = \arctan \left  \frac{y}{x} \right $ ( $\phi$ depends on quadrant)	N/A
<b>Cylindrical</b>	$r = \sqrt{x^2 + y^2}$	$\phi' = \arctan \left  \frac{y}{x} \right $ ( $\phi$ depends on quadrant)	$z = z$
<b>Spherical</b>	$r = \sqrt{x^2 + y^2 + z^2}$	$\phi' = \arctan \left  \frac{y}{x} \right $ ( $\phi$ depends on quadrant)	$\theta = \arccos \left( \frac{z}{\sqrt{x^2 + y^2 + z^2}} \right)$

### 3.3.7 Hyperbolic Functions

#### Hyperbolic Sine

- $\sinh x = \frac{e^x - e^{-x}}{2} = \frac{e^{2x} - 1}{2e^x} = \frac{1 - e^{-2x}}{2e^{-x}}$
- $\sinh x = -i \sin(ix)$

#### Hyperbolic Cosine

- $\cosh x = \frac{e^x + e^{-x}}{2} = \frac{e^{2x} + 1}{2e^x} = \frac{1 + e^{-2x}}{2e^{-x}}$
- $\cosh x = \cos(ix)$

#### Hyperbolic Tangent

- $\tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{e^{2x} - 1}{e^{2x} + 1} = \frac{1 - e^{-2x}}{1 + e^{-2x}}$
- $\tanh x = -i \tan(ix)$

#### Hyperbolic Cotangent

- $\coth x = \frac{1}{\tanh x} = \frac{\cosh x}{\sinh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}} = \frac{e^{2x} + 1}{e^{2x} - 1} = \frac{1 + e^{-2x}}{1 - e^{-2x}}$
- $\coth x = i \cot(ix)$

### Hyperbolic Secant

- $\operatorname{sech} x = \frac{1}{\cosh x} = \frac{2}{e^x + e^{-x}} = \frac{2e^x}{e^{2x} + 1} = \frac{2e^{-x}}{1 + e^{-2x}}$
- $\operatorname{sech} x = \sec(ix)$

### Hyperbolic Cosecant

- $\operatorname{csch} x = \frac{1}{\sinh x} = \frac{2}{e^x - e^{-x}} = \frac{2e^x}{e^{2x} - 1} = \frac{2e^{-x}}{1 - e^{-2x}}$
- $\operatorname{csch} x = i \csc(ix)$

### Identities

- $\cosh^2 x - \sinh^2 x = 1$
- $\sin \theta \cos \theta = \frac{1}{2} \sin(2\theta)$

## 3.4 Trigonometry

### Definitions

- SOH CAH TOA
- $\sin \theta = \frac{\textit{opposite}}{\textit{hypotenuse}}$
- $\cos \theta = \frac{\textit{adjacent}}{\textit{hypotenuse}}$
- $\tan \theta = \frac{\textit{opposite}}{\textit{adjacent}} = \frac{\sin \theta}{\cos \theta}$
- $\cot \theta = \frac{\textit{adjacent}}{\textit{opposite}} = \frac{\cos \theta}{\sin \theta}$
- $\sec \theta = \frac{\textit{hypotenuse}}{\textit{adjacent}}$
- $\csc \theta = \frac{\textit{hypotenuse}}{\textit{opposite}}$

### 3.4.1 Identities

#### Pythagorean Identities

- $\cos^2 \theta + \sin^2 \theta = 1$
- $\tan^2 \theta + 1 = \sec^2 \theta$
- $1 + \cot^2 \theta = \csc^2 \theta$

#### Reciprocals

- $\sin \theta = \frac{1}{\csc \theta}$
- $\cos \theta = \frac{1}{\sec \theta}$
- $\tan \theta = \frac{1}{\cot \theta}$
- $\cot \theta = \frac{1}{\tan \theta}$

- $\sec \theta = \frac{1}{\cos \theta}$
- $\csc \theta = \frac{1}{\sin \theta}$

#### As complex exponentials

- $\sin \theta = \frac{e^{ix} - e^{-ix}}{2i}$
- $\cos \theta = \frac{e^{ix} + e^{-ix}}{2}$
- $\tan \theta = \frac{e^{ix} - e^{-ix}}{i(e^{ix} + e^{-ix})}$
- $\cot \theta = \frac{i(e^{ix} + e^{-ix})}{e^{ix} - e^{-ix}}$
- $\sec \theta = \frac{2}{e^{ix} + e^{-ix}}$
- $\csc \theta = \frac{2i}{e^{ix} - e^{-ix}}$

#### Symmetries

- $\sin(-\theta) = -\sin \theta$
- $\cos(-\theta) = \cos \theta$
- $\tan(-\theta) = -\tan \theta$
- $\csc(-\theta) = -\csc \theta$
- $\sec(-\theta) = \sec \theta$
- $\cot(-\theta) = -\cot \theta$
  
- $\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$
- $\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$
- $\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta$
- $\csc\left(\frac{\pi}{2} - \theta\right) = \sec \theta$
- $\sec\left(\frac{\pi}{2} - \theta\right) = \csc \theta$
- $\cot\left(\frac{\pi}{2} - \theta\right) = \tan \theta$
  
- $\sin(\pi - \theta) = \sin \theta$
- $\cos(\pi - \theta) = -\cos \theta$
- $\tan(\pi - \theta) = -\tan \theta$
- $\csc(\pi - \theta) = \csc \theta$
- $\sec(\pi - \theta) = -\sec \theta$
- $\cot(\pi - \theta) = -\cot \theta$

### Angle sum and difference formulae

- $\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$
- $\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$
- $\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$

### Half-angle formulae

- $\sin^2\left(\frac{\theta}{2}\right) = \frac{1 - \cos \theta}{2}$
- $\cos^2\left(\frac{\theta}{2}\right) = \frac{1 + \cos \theta}{2}$
- $\tan^2\left(\frac{\theta}{2}\right) = \frac{1 - \cos \theta}{1 + \cos \theta}$
- $\tan\left(\frac{\theta}{2}\right) = \frac{\tan \theta}{1 + \sec \theta} = \frac{\sin \theta}{1 + \cos \theta} = \frac{1 - \cos \theta}{\sin \theta} = \csc \theta - \cot \theta$

### Double-angle formulae

- $\cos(2\theta) = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta = \cos^2 \theta - \sin^2 \theta$
- $\sin(2\theta) = 2 \sin \theta \cos \theta$
- $\tan(2\theta) = \frac{2 \tan \theta}{1 - \tan^2 \theta}$

### Sum to Product

- $\sin \alpha + \sin \beta = 2 \sin\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha - \beta}{2}\right)$
- $\sin \alpha - \sin \beta = 2 \cos\left(\frac{\alpha + \beta}{2}\right) \sin\left(\frac{\alpha - \beta}{2}\right)$
- $\cos \alpha + \cos \beta = 2 \cos\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha - \beta}{2}\right)$
- $\cos \alpha - \cos \beta = -2 \sin\left(\frac{\alpha + \beta}{2}\right) \sin\left(\frac{\alpha - \beta}{2}\right)$

### Product to Sum

- $\sin \alpha \sin \beta = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$
- $\cos \alpha \cos \beta = \frac{1}{2}[\cos(\alpha - \beta) + \cos(\alpha + \beta)]$
- $\sin \alpha \cos \beta = \frac{1}{2}[\sin(\alpha + \beta) + \cos(\alpha - \beta)]$
- $\cos \alpha \sin \beta = \frac{1}{2}[\sin(\alpha + \beta) - \sin(\alpha - \beta)]$

### Law of Sines

- $\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$

### Law of Cosines

- $a^2 = b^2 + c^2 - 2bc \cos \alpha$
- $b^2 = a^2 + c^2 - 2ac \cos \beta$
- $c^2 = a^2 + b^2 - 2ab \cos \gamma$

### Law of Tangents

- $\frac{a-b}{a+b} = \frac{\tan(\frac{1}{2}[\alpha-\beta])}{\tan(\frac{1}{2}[\alpha+\beta])}$
- $\frac{b-c}{b+c} = \frac{\tan(\frac{1}{2}[\beta-\gamma])}{\tan(\frac{1}{2}[\beta+\gamma])}$
- $\frac{a-c}{a+c} = \frac{\tan(\frac{1}{2}[\alpha-\gamma])}{\tan(\frac{1}{2}[\alpha+\gamma])}$

### Mollweide's Formula

- $\frac{a+b}{c} = \frac{\cos(\frac{1}{2}[\alpha-\beta])}{\sin(\frac{1}{2}\gamma)}$

### Small-angle approximations

- $\sin \theta \approx \theta$
- $\cos \theta \approx 1 - \frac{\theta^2}{2}$
- $\tan \theta \approx \theta$

### Other identities

- $\sin \theta \cos \theta = \frac{1}{2} \sin(2\theta)$
- $\cos^2 \theta = \frac{1}{2}(\cos(2\theta) + 1)$

### Averages

- $\overline{\sin x} = \overline{\cos x} = 0$
- $\overline{\sin^2 x} = \overline{\cos^2 x} = \frac{1}{2}$

### Table of Identities

In terms of...	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\sec \theta$	$\cot \theta$	$\csc \theta$
$\sin \theta =$	$\sin \theta$	$\pm\sqrt{1 - \cos^2 \theta}$	$\pm\frac{\tan \theta}{\sqrt{1 + \tan^2 \theta}}$	$\pm\frac{\sqrt{\sec^2 \theta - 1}}{\sec \theta}$	$\pm\frac{1}{\sqrt{1 + \cot^2 \theta}}$	$\frac{1}{\csc \theta}$
$\cos \theta =$	$\pm\sqrt{1 - \sin^2 \theta}$	$\cos \theta$	$\pm\frac{1}{\sqrt{1 + \tan^2 \theta}}$	$\frac{1}{\sec \theta}$	$\pm\frac{\cot \theta}{\sqrt{1 + \cot^2 \theta}}$	$\pm\frac{\sqrt{\csc^2 - 1}}{\csc \theta}$
$\tan \theta =$	$\pm\frac{\sin \theta}{\sqrt{1 - \sin^2 \theta}}$	$\pm\frac{\sqrt{1 - \cos^2 \theta}}{\cos \theta}$	$\tan \theta$	$\pm\sqrt{\sec^2 \theta - 1}$	$\frac{1}{\cot \theta}$	$\pm\frac{1}{\sqrt{\csc^2 \theta - 1}}$
$\sec \theta =$	$\pm\frac{1}{\sqrt{1 - \sin^2 \theta}}$	$\frac{1}{\cos \theta}$	$\pm\sqrt{1 + \tan^2 \theta}$	$\sec \theta$	$\pm\frac{\sqrt{1 + \cot^2 \theta}}{\cot \theta}$	$\pm\frac{\csc \theta}{\sqrt{\csc^2 - 1}}$
$\cot \theta =$	$\pm\frac{\sqrt{1 - \sin^2 \theta}}{\sin \theta}$	$\pm\frac{\cos \theta}{\pm\sqrt{1 - \cos^2 \theta}}$	$\frac{1}{\tan \theta}$	$\pm\frac{1}{\sqrt{\sec^2 \theta - 1}}$	$\cot \theta$	$\pm\sqrt{\csc^2 \theta - 1}$
$\csc \theta =$	$\frac{1}{\sin \theta}$	$\pm\frac{1}{\sqrt{1 - \cos^2 \theta}}$	$\pm\frac{\sqrt{1 + \tan^2 \theta}}{\tan \theta}$	$\pm\frac{\sec \theta}{\sqrt{\sec^2 \theta - 1}}$	$\pm\sqrt{1 + \cot^2 \theta}$	$\csc \theta$

## 3.5 Calculus

### 3.5.1 Limits

### 3.5.2 Properties

- $\lim_{x \rightarrow a} cf(x) = c \lim_{x \rightarrow a} f(x)$
- $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \quad \lim_{x \rightarrow a} g(x) \neq 0$
- $\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$

#### Useful Limits

- $\lim_{x \rightarrow \infty} e^x = \infty$
- $\lim_{x \rightarrow -\infty} e^x = 0$
- $\lim_{x \rightarrow \infty} \ln(x) = \infty$
- $\lim_{x \rightarrow 0^-} \ln(x) = -\infty$
- $\lim_{x \rightarrow 0} x \log x = 0$

#### L'Hôpital's rule

- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

#### Squeeze principle

- For  $g(x) \leq f(x) \leq h(x)$  and  $\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L$ :  

$$\lim_{x \rightarrow a} f(x) = L$$

### 3.5.3 Differentiation

#### First Principles

- $\frac{d}{dx} f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

#### Nature of derivatives

Derivative	Function
$f'x > 0$	Increasing
$f'x = 0$	Stationary
$f'x < 0$	Decreasing

Second Derivative	Function	Stationary Points [ $f'x = 0$ ]
$f''x > 0$	Concave up	Local Minimum
$f'x = 0$	No information	Inflection Point
$f''x < 0$	Concave down	Local Maximum



### Product Rule

- $(uv)' = uv' + vu'$
- $\frac{d}{dx}f(x)g(x) = f(x)g'(x) + f'(x)g(x)$

### Quotient Rule

- $\left(\frac{u}{v}\right)' = \frac{vu' - uv'}{v^2}$
- $\frac{d}{dx} \frac{f(x)}{g(x)} = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$

### Chain Rule

- $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
- $\frac{d}{dx}f(g(x)) = f'(g(x))g'(x)$

### Useful Derivatives

- $\frac{d}{dx}x^n = nx^{n-1}$
- $\frac{d}{dx}a^x = a^x \ln(a)$
- $\frac{d}{dx}e^x = e^x$
- $\frac{d}{dx} \ln x = \frac{1}{x}, x > 0$
- $\frac{d}{dx} \ln |x| = \frac{1}{x}, x \neq 0$
- $\frac{d}{dx} \ln(f(x)) = \frac{f'(x)}{f(x)}$
- $\frac{d}{dx} \log_b x = \frac{1}{x \ln(b)}, x > 0$
- $\frac{d}{dx} \sin x = \cos x$
- $\frac{d}{dx} \cos x = -\sin x$
- $\frac{d}{dx} \tan x = \sec^2 x$
- $\frac{d}{dx} \sec x = \sec x \tan x$
- $\frac{d}{dx} \csc x = -\csc x \cot x$
- $\frac{d}{dx} \cot x = -\csc^2 x$
- $\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$
- $\frac{d}{dx} \cos^{-1} x = \frac{1}{-\sqrt{1-x^2}}$
- $\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$

### 3.5.4 Partial Differentiation

#### First Principles

- $$\frac{\partial}{\partial x} f(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}$$

#### Jacobian Matrix

- $$D\vec{f}(\vec{a}) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1}(\vec{a}) & \frac{\partial f_1}{\partial x_2}(\vec{a}) & \cdots & \frac{\partial f_1}{\partial x_n}(\vec{a}) \\ \frac{\partial f_2}{\partial x_1}(\vec{a}) & \frac{\partial f_2}{\partial x_2}(\vec{a}) & \cdots & \frac{\partial f_2}{\partial x_n}(\vec{a}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1}(\vec{a}) & \frac{\partial f_m}{\partial x_2}(\vec{a}) & \cdots & \frac{\partial f_m}{\partial x_n}(\vec{a}) \end{bmatrix}$$

#### Definition of differentiability of a multivariable function

- $$\lim_{\vec{x} \rightarrow \vec{a}} \frac{\|f(\vec{x}) - f(\vec{a}) - Df(\vec{a}) \cdot (\vec{x} - \vec{a})\|}{\|\vec{x} - \vec{a}\|} = 0$$

### 3.5.5 The Differential

- $$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy + \frac{\partial F}{\partial z} dz$$

### 3.5.6 Line Element

- $$dS^2 = dx^2 + dy^2 + dz^2$$

### 3.5.7 Integration

#### Properties

- $$\int f(x) \pm g(x) dx = \left( \int f(x) dx \right) \pm \left( \int g(x) dx \right)$$
- $$\int_a^a f(x) dx = 0$$
- $$\int_a^b f(x) dx = - \int_b^a f(x) dx$$
- $$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$
- If  $f(x) \geq g(x)$  over  $[a, b]$ ,  $\int_a^b f(x) dx \geq \int_a^b g(x) dx$
- If  $f(x) \geq 0$  over  $[a, b]$ ,  $\int_a^b f(x) dx > 0$
- If  $m \leq f(x) \leq M$  over  $[a, b]$ ,  $m(b - a) \leq \int_a^b f(x) dx \leq M(b - a)$

### Integration by Parts

- $\int u'v = uv - \int uv'$
- $\int f'(x)g(x) dx = f(x)g(x) - \int f(x)g'(x) dx$
- $\int_a^b f'(x)g(x) dx = [f(x)g(x)]_a^b - \int_a^b f(x)g'(x) dx$

### Integration by Substitution

- $u = g(x); dx = g'(x) dx; \int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(x) dx$

### Approximations

*Trapezoid rule*

- $\int_a^b f(x) dx \approx \frac{\Delta x}{2}[f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)]$

*Simpson's rule*

- $\int_a^b f(x) dx \approx \frac{\Delta x}{3}[f(x_0) + 4f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$

### Useful Indefinite Integrals

- $\int k dx = kx + C$
- $\int \log_b x dx = x(\log_b x - \log_b(e)) + C = x(\log_b x - \frac{1}{\ln b}) + C$
- $\int \ln x dx = x(\ln x - 1) + C$
- $\int e^x dx = e^x + C$
- $\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$
- $\int \frac{1}{x} dx = \ln|x| + C$
- $\int \frac{1}{ax+b} dx = \frac{1}{a} \ln|ax+b| + C$
- $\int \sin x dx = -\cos x + C$
- $\int \cos x dx = \sin x + C$
- $\int \tan x dx = \ln|\sec x| + C$
- $\int \sec x dx = \ln|\sec x + \tan x| + C$
- $\int \sec^2 x dx = \tan x + C$
- $\int \csc^2 x dx = -\cot x + C$
- $\int \sec x \tan x dx = \sec x + C$
- $\int \csc x \cot x dx = -\csc x + C$
- $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1}(\frac{x}{a}) + C$
- $\int \frac{1}{\sqrt{a^2 + x^2}} dx = \sin^{-1}(\frac{x}{a}) + C$

### Useful Definite Integrals

- $\int_0^{\infty} \frac{x^3}{e^x - 1} dx = \frac{\pi^4}{15}$
- $\int_0^{\infty} x^n e^{-ax} dx = \frac{n!}{a^{n+1}}$

### 3.5.8 Vector Calculus

#### Vector derivative

- $\text{grad}(f) = \vec{\nabla} f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z}$
- $\int_a^b (\vec{\nabla} f) \cdot d\vec{l} = f(a) - f(b)$

#### The Laplacian

- $\delta f = \nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$

#### Divergence

##### *In Cartesian Coordinates*

- $\text{div}(f) = \vec{\nabla} \cdot \vec{f} = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} + \frac{\partial f_z}{\partial z}$

##### *In Cylindrical Coordinates*

- $\vec{\nabla} \cdot \vec{f} = \frac{1}{r} \frac{\partial}{\partial r} (r f_r) + \frac{1}{r} \frac{\partial f_\phi}{\partial \phi} + \frac{\partial f_z}{\partial z}$

##### *In Spherical Coordinates*

- $\vec{\nabla} \cdot \vec{f} = \frac{1}{r} \frac{\partial}{\partial r} (r^2 f_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (f_\theta \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial f_\phi}{\partial \phi}$

#### Curl

##### *In Cartesian Coordinates*

- $\text{curl}(f) = \vec{\nabla} \times \vec{f} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_x & f_y & f_z \end{vmatrix} = \left( \frac{\partial f_z}{\partial y} - \frac{\partial f_y}{\partial z} \right) \hat{x} + \left( \frac{\partial f_x}{\partial z} - \frac{\partial f_z}{\partial x} \right) \hat{y} + \left( \frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) \hat{z}$

##### *In Cylindrical Coordinates*

- $\vec{\nabla} \times \vec{f} = \left( \frac{1}{r} \frac{\partial f_z}{\partial \phi} - \frac{\partial f_\phi}{\partial z} \right) \hat{r} + \left( \frac{\partial f_r}{\partial z} - \frac{\partial f_z}{\partial r} \right) \hat{\phi} + \frac{1}{r} \left( \frac{\partial}{\partial r} (r f_\phi) - \frac{\partial f_r}{\partial \phi} \right) \hat{z}$

##### *In Spherical Coordinates*

- $\vec{\nabla} \times \vec{f} = \frac{1}{r \sin \theta} \left( \frac{\partial}{\partial \theta} (f_\phi \sin \theta) - \frac{\partial f_\theta}{\partial \phi} \right) \hat{r} + \frac{1}{r} \left( \frac{1}{\sin \theta} \frac{\partial f_r}{\partial \phi} - \frac{\partial}{\partial r} (r f_\phi) \right) + \frac{1}{r} \left( \frac{\partial}{\partial r} (r f_\theta) - \frac{\partial f_r}{\partial \theta} \right)$

### Vector Second Derivatives

- $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{v}) = 0$
- $\vec{\nabla}(\vec{\nabla}(\vec{\nabla} \cdot \vec{f}))$

*Vector Laplacian*

- $\vec{\nabla}^2 \vec{f} = \vec{\nabla}(\vec{\nabla} \cdot \vec{f}) - \vec{\nabla} \times (\vec{\nabla} \times \vec{f})$

### Stokes' Theorem

- $\iint_S (\vec{\nabla} \times \vec{f}) \cdot d\vec{a} = \oint_B \vec{v} \cdot d\vec{l}$

### Divergence Theorem

- $\iiint_V (\vec{\nabla} \cdot \vec{f}) dV = \oint_S \vec{v} \cdot d\vec{a}$

### 3.5.9 Dirac Delta Function

- $\delta(x) = \begin{cases} 0 & x \neq 0 \\ \infty & x = 0 \end{cases}$
- $\delta(x - a) = \begin{cases} 0 & x \neq a \\ \infty & x = a \end{cases}$
- $\int_{-\infty}^{\infty} \delta(x) dx = 1$
- $f(x)\delta(x) = f(0)\delta(x)$

### 3.5.10 Approximations

- $f(x + \Delta x) \approx f(x) + \Delta x f'(x)$

# Chapter 4

## Statistics

### 4.1 Variance

- $var(x) = \langle (x - \langle x \rangle)^2 \rangle = \langle x^2 \rangle - \langle x \rangle^2$

### 4.2 Standard Deviation

- $\sigma(x) = \sqrt{var(x)} = \sqrt{\langle x^2 \rangle - \langle x \rangle^2}$

#### 4.2.1 Population Standard Deviation

- $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$

#### 4.2.2 Sample Standard Deviation

- $\sigma_{sample} = \sqrt{\frac{1}{N-1} \sum_{i=1}^N (x_i - \bar{x})^2}$

### 4.3 Residual Sum of Squares

- $RSS = \sum_{i=1}^n (y_i - \hat{y}_i)^2$

### 4.4 Mean Squared Error

- $MSE = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{f}(x_i))^2 = \frac{1}{n} RSS = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$

### 4.5 Residual Standard Error

- $RSE = \sqrt{\frac{1}{n-2} RSS} = \sqrt{\frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$

### 4.6 Correlation

- $Cor(X, Y) = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$

# Appendix A

## Values

### A.1 Physics

#### A.1.1 Physical Constants

$c$ : Speed of light

$$\bullet = \frac{1}{\sqrt{\varepsilon_0 \mu_0}} = 299\,792\,458 \text{ m} \cdot \text{s}^{-1} \approx 3 \times 10^8 \text{ m} \cdot \text{s}^{-1}$$

$G$ : Universal gravitational constant

$$\bullet = 6.67408(31) \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2} \approx 6.67 \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}$$

$g$ : Average acceleration due to gravity at sea level on Earth

$$\bullet = 9.80665 \text{ m} \cdot \text{s}^{-2} \approx 9.8 \text{ m} \cdot \text{s}^{-2}$$

$h$ : Planck constant

$$\bullet = 6.626\,070\,040 \times 10^{-34} \text{ J} \cdot \text{s} \approx 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$$

$\hbar$ : Reduced Planck constant

$$\bullet = \frac{h}{2\pi} = 1.054\,571\,726 \times 10^{-34} \text{ J} \cdot \text{s} \approx 1.055 \times 10^{-34} \text{ J} \cdot \text{s}$$

$k_B$ : Boltzmann constant

$$\bullet = 1.3\,806\,488 \times 10^{-23} \text{ J} \cdot \text{K}^{-1} \approx 1.38 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$$

$k_e$ : Coulomb's constant

$$\bullet = \frac{1}{4\pi\varepsilon_0} = 8.987\,551\,787 \times 10^9 \text{ N} \cdot \text{m} \cdot \text{C}^{-2} \approx 9 \times 10^9 \text{ N} \cdot \text{m} \cdot \text{C}^{-2}$$

$N_A$ : Avogadro constant

$$\bullet = 6.022\,140\,857(74) \times 10^{23} \text{ mol}^{-1} \approx 6.022 \times 10^{23} \text{ mol}^{-1}$$

$\varepsilon_0$ : Vacuum permittivity

$$\bullet = \frac{1}{\mu_0 c^2} = 8.854\,187\,817 \times 10^{-12} \text{ F} \cdot \text{m}^{-1} \approx 8.85 \times 10^{-12} \text{ F} \cdot \text{m}^{-1}$$

$\mu_0$ : Vacuum permeability

$$\bullet = 4\pi \times 10^{-7} \text{ N} \cdot \text{A}^{-2} = \frac{1}{\varepsilon_0 c^2} \approx 1.257 \times 10^{-6} \text{ N} \cdot \text{A}^{-2}$$

### A.1.2 Useful Quantities

Density of air ( $\rho_A$ ):  $1.2922 \text{ kg} \cdot \text{m}^{-3}$

Density of water ( $\rho_w$ ):  $10^3 \text{ kg} \cdot \text{m}^{-3}$

Mass of an electron ( $m_e$ ):  $9.10938291 \times 10^{-31} \text{ kg} \approx 9 \times 10^{-31} \text{ kg}$

Mass of a neutron ( $m_n$ ):  $1.674927351 \times 10^{-27} \text{ kg} \approx 1.675 \times 10^{-27} \text{ kg}$

Mass of a proton ( $m_p$ ):  $1.672621777 \times 10^{-27} \text{ kg} \approx 1.672 \times 10^{-27} \text{ kg}$

Speed of sound in air:  $343.2 \text{ m} \cdot \text{s}^{-1}$

Specific heat capacity of water:  $4.186 \times 10^3 \text{ J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$

## A.2 Astronomy

### A.2.1 Useful Quantities

Surface Temperature of the Sun: ( $T_\odot$ ):  $= 5778 \text{ K} = 5505 \text{ }^\circ\text{C}$

#### Planetary Properties

Body	Mass	Average Radius	Semi-major axis	Eccentricity	Orbital period
Mercury ♿	$3.3011 \times 10^{23} \text{ kg}$ $0.055 M_\oplus$ $1.66 \times 10^{-7} M_\odot$ $1.74 \times 10^{-4} M_{\text{Jup}}$	$2.4397 \times 10^6 \text{ m}$ $0.3829 R_\oplus$	$5.790905 \times 10^{10} \text{ m}$ $0.387098 \text{ AU}$	0.205630	0.240856 yr
Venus ♀	$4.8675 \times 10^{24} \text{ kg}$ $0.815 M_\oplus$ $2.447 \times 10^{-6} M_\odot$ $2.56 \times 10^{-3} M_{\text{Jup}}$	$6.0518 \times 10^6 \text{ m}$ $0.9499 R_\oplus$	$1.08208000 \times 10^{11} \text{ m}$ $0.723332 \text{ AU}$	0.006772	0.615198 yr
Earth ☉	$5.97237 \times 10^{24} \text{ kg}$ $1 M_\oplus$ $3.003 \times 10^{-6} M_\odot$ $2.67 \times 10^{-3} M_{\text{Jup}}$	$6.3710 \times 10^6 \text{ m}$ $1 R_\oplus$	$1.49598023 \times 10^{11} \text{ m}$ $1.000001 \text{ AU}$	0.0167086	1.000017 yr
Mars ♂	$6.4171 \times 10^{23} \text{ kg}$ $0.107 M_\oplus$ $3.226 \times 10^{-7} M_\odot$ $3.38 \times 10^{-4} M_{\text{Jup}}$	$3.3895 \times 10^6 \text{ m}$ $0.53 R_\oplus$	$2.27939200 \times 10^{11} \text{ m}$ $1.523679 \text{ AU}$	0.0934	1.88082 yr
Jupiter ♃	$1.8982 \times 10^{27} \text{ kg}$ $317.8 M_\oplus$ $9.55 \times 10^{-4} M_\odot$ $1 M_{\text{Jup}}$	$6.9911 \times 10^7 \text{ m}$ $10.97 R_\oplus$	$7.4052 \times 10^{11} \text{ m}$ $5.2044 \text{ AU}$	0.0489	11.862 yr
Saturn ♄	$5.6834 \times 10^{26} \text{ kg}$ $95.159 M_\oplus$ $2.86 \times 10^{-4} M_\odot$ $0.299 M_{\text{Jup}}$	$5.8232 \times 10^7 \text{ m}$ $9.14 R_\oplus$	$1.43353 \times 10^{12} \text{ m}$ $9.5826 \text{ AU}$	0.0565	29.4571 yr
Uranus ♅	$8.68 \times 10^{25} \text{ kg}$ $14.536 M_\oplus$ $4.36 \times 10^{-5} M_\odot$ $0.046 M_{\text{Jup}}$	$2.5362 \times 10^7 \text{ m}$ $3.98 R_\oplus$	$2.87504 \times 10^{12} \text{ m}$ $19.2184 \text{ AU}$	0.046381	84.0205 yr
Neptune ♆	$1.0243 \times 10^{26} \text{ kg}$ $17.147 M_\oplus$ $5.15 \times 10^{-5} M_\odot$ $0.054 M_{\text{Jup}}$	$2.4622 \times 10^7 \text{ m}$ $3.86 R_\oplus$	$4.5 \times 10^{12} \text{ m}$ $30.11 \text{ AU}$	0.009456	164.8 yr



### A.3 Mathematics

**Euler's number ( $e$ ):**  $\sum_{n=0}^{\infty} \frac{1}{n!} = \frac{1}{1} + \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \dots$   
= 2.71828182845904523536028747135266249775724709369995...  
 $\approx 2.7182 \quad \approx 2.718$

**Pi ( $\pi$ ):**  $\frac{C}{d} = \frac{C}{2r}$   
= 3.14159265358979323846264338327950288419716939937510...  
 $\approx 3.14159 \quad \approx 3.142$

# Appendix B

## Units of Measurement

### B.1 Natural Units

Handy when you're dealing with small things.

**Charge:** elementary charge ( $e$ )

- The electric charge of a proton.
- $= 1.602\,176\,565 \times 10^{-19} \text{ C} \approx 1.6 \times 10^{-19} \text{ C}$

**Energy:** electron volt ( $eV$ )

- The work done to move an electron across one volt of potential.
- $= e \cdot V = 1.602\,176\,565 \times 10^{-19} \text{ J} \approx 1.6 \times 10^{-19} \text{ J}$

### B.2 SI System

Universally acknowledged as the best system of units.

#### B.2.1 Base Units

**Amount of Substance:** mole ( $mol$ )

- The amount of substance of a system which contains as many elementary entities as there are atoms in 0.012  $kg$  of carbon-12.
- $= 6.022\,140\,857 \times 10^{23}$

**Electric Current:** ampere ( $A$ )

- The constant current which, if maintained in two straight parallel conductors of infinite length, of negligible circular cross-section, and placed 1 m apart in vacuum, would produce between these conductors a force equal to  $2 \times 10^{-7} \text{ N} \cdot \text{m}^{-1}$  of length.
- $= \text{C} \cdot \text{s}^{-1}$

**Length:** metre ( $m$ )

- The distance traveled by light in vacuum in  $\frac{1}{299\,792\,458} \text{ s}$
- $= 3.2808 \text{ ft}$

**Luminous intensity:** candela ( $cd$ )

- The luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency  $5.4 \times 10^{14} \text{ Hz}$  and that has a radiant intensity in that direction of  $\frac{1}{683}$  watt per steradian.

**Mass:** kilogram ( $kg$ )

- $= 2.205lb$

**Thermodynamic Temperature:** kelvin ( $K$ )

- $\frac{1}{273.16}$  of the thermodynamic temperature of the triple point of water.
- $= T[^\circ C] + 273.15$

**Time:** second ( $s$ )

- The duration of 9 192 631 770 periods of rotation corresponding to the two hyperfine levels of the ground-state of the caesium-133 atom.

## B.2.2 Derived Units

**Angle:** radian ( $rad$ )

- A full circle divided by  $2\pi$ .
- $= m \cdot m^{-1} = \frac{180}{\pi} = 206265 \text{ arcsecs} \approx 57.3^\circ$

**Electric Charge:** coulomb ( $C$ )

- $= A \cdot s = 6.242 \times 10^{18} e$

**Electrical capacitance:** farad ( $F$ )

- $= m^{-2} \cdot kg^{-1} \cdot s^4 \cdot A^2$

**Electrical conductance:** siemens ( $S$ )

- $= A \cdot V^{-1} = kg^{-1} \cdot m^{-2} \cdot s^3 \cdot A^2$

**Electrical inductance:** henry ( $H$ )

- $= Wb \cdot A^{-1} = kg \cdot m^2 \cdot s^{-2} \cdot A^{-2}$

**Electrical potential difference / Voltage:** volt ( $V$ )

- $= W \cdot A^{-1} = kg \cdot m^2 \cdot s^{-3} \cdot A^{-1}$

**Electrical resistance:** ohm ( $\Omega$ )

- $= V \cdot A^{-1} = kg \cdot m^2 \cdot s^{-3} \cdot A^{-2}$

**Energy:** joule ( $J$ )

- $= N \cdot m = kg \cdot m^2 \cdot s^{-2}$

**Force:** newton ( $N$ )

- $= kg \cdot m \cdot s^{-2} = 0.224809 \text{ lbf}$

**Frequency:** hertz ( $Hz$ )

- $= s^{-1}$

**Illuminance:** lux ( $lx$ )

- $= lm \cdot m^{-2} = m^{-2} \cdot cd$

**Luminous flux:** lumen ( $lm$ )

- $= cd \cdot sr = cd$

**Magnetic flux:** weber ( $Wb$ )

- $= V \cdot s = kg \cdot m^2 \cdot s^{-2} \cdot A^{-1}$

**Magnetic flux density:** tesla ( $T$ )

- $= kg \cdot s^{-2} \cdot A^{-1}$

**Power:** watt ( $W$ )

- $= J \cdot s = kg \cdot m^2 \cdot s^{-3}$

**Pressure:** pascal ( $Pa$ )

- $= N \cdot m^{-2} = kg \cdot m^{-1} \cdot s^{-2}$

**Radioactivity:** becquerel ( $\Omega$ )

- Decays per second
- $= s^{-1}$

**Solid angle:** steradian ( $sr$ )

- $= m^2 \cdot m^{-2}$

**Temperature:** degree Celcius ( $^{\circ}C$ )

- $T[C] = T[K] - 273.15$

### B.3 CGS (centimetres-grams-seconds)

Commonly used in astronomy, to everyone's disappointment.

**Acceleration:** gal ( $Gal$ )

- $= cm \cdot s^{-2} = 10^{-2} m \cdot s^{-2}$

**Energy:** erg ( $erg$ )

- $= g \cdot cm^2 \cdot s^{-2} = 10^{-7} J$

**Force:** dyne ( $dyn$ )

- $= g \cdot cm \cdot s^{-2} = 10^{-5} N$

**Length:** centimetre ( $cm$ )

- $= 0.01 m$

**Mass:** gram ( $g$ )

- $= 10^{-3} kg$

**Power:** erg per second ( $erg/s$ )

- $= g \cdot cm^2 \cdot s^{-2} = 10^{-7} W$

**Pressure:** barye ( $Ba$ )

- $= g \cdot cm^{-1} \cdot s^{-2} = 10^{-1} Pa$

**Time:** second ( $s$ )

**Velocity:** centimetre per second ( $cm/s$ )

- $= 10^{-2} m \cdot s^{-1}$

**Viscosity (dynamic):** poise ( $P$ )

- $= g \cdot cm^{-1} s^{-1} = 10^{-1} Pa \cdot s$

**Viscosity (kinematic):** stokes ( $St$ )

- $= g \cdot cm^2 s^{-1} = 10^{-4} m^2 \cdot s^{-1}$

**Wavenumber:** kayser ( $K$ )

- $= cm^{-1} = 100 m^{-1}$

## B.4 Astronomy units

### B.4.1 Astronomical system

**Distance:** astronomical unit ( $AU$ )

- Roughly the distance from the Earth to the Sun.
- $= 1.4960 \times 10^{11} m = 4.8481 \times 10^{-6} pc = 1.5813 \times 10^{-5} ly$

**Mass:** solar mass ( $M_{\odot}$ )

- $= 1.98855 \times 10^{30} kg \approx 2 \times 10^{30} kg = 1048 M_{\oplus} = 332\,950 M_{\odot}$

**Time:** Day

- $= 86\,400s$

#### Complimentary units

**Distance:** Solar radius ( $R_{\odot}$ )

- $= 6.957 \times 10^8 m = 695\,700 km \approx 7 \times 10^8 m$

**Distance:** parsec ( $pc$ )

- The distance at which the parallax of an object over the course of the Earth's orbit is one arcsec.
- $= 3.0857 \times 10^{16} m = 2.0626 \times 10^5 AU = 3.26156 ly$

**Distance:** light year ( $ly$ )

- The distance travelled by light in a vacuum in a year.
- $= 9.4607 \times 10^{15} m = 6.3241 \times 10^4 AU = 0.3066 pc$

**Mass:** Earth mass ( $M_{\oplus}$ )

- $= 5.9722 \times 10^{24} kg \approx 6 \times 10^{27} kg$

**Mass:** Jupiter mass ( $M_{\text{Jup}}$ )

- $= 1.898 \times 10^{27} kg \approx 1.9 \times 10^{27} kg$

**Specific Flux:** Jansky ( $Jy$ )

- $= 10^{-26} W \cdot m^{-2} \cdot Hz^{-1}$

### B.4.2 Equatorial Coordinate System

**Right Ascension** ( $\alpha$ )

**Hour** ( $^h$ ):  $\frac{1}{24} circle = 15^\circ$

**Minute** ( $^m$ ):  $\frac{1}{60}^h = \frac{1}{1440} circle = 15'$

**Second** ( $^s$ ):  $\frac{1}{60}^m = \frac{1}{3600}^h = \frac{1}{86400} circle = 15''$

**Declination** ( $\delta$ )

Declination is measured using normal degrees (see *Degrees of Angle*) from the equator.

## B.5 United States customary units (aka Imperial Units)

### B.5.1 Length

$$\text{Point } (p): = \frac{127}{360} \text{ mm}$$

$$\text{Pica } (P/): = 12 p = \frac{127}{30} \text{ mm}$$

$$\text{Inch } (in \text{ or } \text{''}): = 6 P/ = 25.4 \text{ mm}$$

$$\text{Foot } (ft \text{ or } \text{'}): = 12 in = 0.3048 \text{ m}$$

$$\text{Yard } (yd): = 3 ft = 0.9144 \text{ m}$$

$$\text{Mile } (Mi): = 5280 ft = 1760 yd = 1.609344 \text{ km}$$

## B.6 Degrees of Angle

$$\text{Degree } (^\circ): \frac{1}{360} \text{ circle} = \frac{\pi}{180} \text{ rad} \approx 0.0174532925199433 \text{ rad}$$

$$\text{Minute of arc } (arcmin \text{ or } \text{'}): \frac{1}{60} ^\circ = \frac{1}{21600} \text{ circle} = \frac{\pi}{10800} \text{ rad}$$

$$\text{Second of arc } (arcsec \text{ or } \text{''}): \frac{1}{60} arcmin = \frac{1}{3600} ^\circ = \frac{1}{206265} \text{ circle} = \frac{\pi}{648000} \text{ rad}$$

## B.7 Miscellaneous Units

### B.7.1 Pressure

$$\text{Bar } (bar): = 10^5 \text{ Pa} \approx 0.9869 \text{ atm}$$

$$\text{Atmosphere } (atm): = 101325 \text{ Pa} = 1.01325 \text{ bar}$$

$$\text{Torr } (torr): = \frac{1}{760} \text{ atm} = \frac{101325}{760} \text{ Pa} \approx 133.3224 \text{ Pa}$$

## B.8 Prefixes

$$\text{atto } (a) = \times 10^{-18}$$

$$\text{femto } (f) = \times 10^{-15}$$

$$\text{pico } (p) = \times 10^{-12}$$

$$\text{nano } (n) = \times 10^{-9}$$

$$\text{micro } (\mu) = \times 10^{-6}$$

$$\text{milli } (m) = \times 10^{-3}$$

$$\text{centi } (c) = \times 10^{-2}$$

$$\text{deca } (da) = \times 10^1$$

$$\text{hecto } (h) = \times 10^2$$

$$\text{kilo } (k) = \times 10^3$$

$$\text{mega } (M) = \times 10^6$$

$$\text{giga } (G) = \times 10^9$$

$$\text{tera } (T) = \times 10^{12}$$


**peta** ( $P$ ) =  $\times 10^{15}$

**exa** ( $E$ ) =  $\times 10^{18}$

# Appendix C



## Mathematical Stuff

### C.1 Trigonometric Values



rad	°	sin	cos	tan
0	0	0	1	0
$\pi/6$	30	1/2	$\sqrt{3}/2$	$1/\sqrt{3}$
$\pi/4$	45	$1/\sqrt{2}$	$1/\sqrt{2}$	1
$\pi/3$	60	$\sqrt{3}/2$	1/2	$\sqrt{3}$
$\pi/2$	90	1	0	$\pm\infty$
$2\pi/3$	120	$\sqrt{3}/2$	-1/2	$-\sqrt{3}$
$3\pi/4$	135	$1/\sqrt{2}$	$-1/\sqrt{2}$	-1
$5\pi/6$	150	1/2	$-\sqrt{3}/2$	$-1/\sqrt{3}$
$\pi$	180	0	-1	0
$7\pi/6$	210	-1/2	$\sqrt{3}/2$	$1/\sqrt{3}$
$5\pi/4$	225	$-1/\sqrt{2}$	$-1/\sqrt{2}$	1
$4\pi/3$	240	$-\sqrt{3}/2$	-1/2	$\sqrt{3}$
$3\pi/2$	270	-1	0	$\pm\infty$
$5\pi/3$	300	$-\sqrt{3}/2$	1/2	$-\sqrt{3}$
$7\pi/4$	315	$-1/\sqrt{2}$	$1/\sqrt{2}$	-1
$11\pi/6$	330	-1/2	$\sqrt{3}/2$	$-1/\sqrt{3}$
$2\pi$	360	0	1	0

### C.1.1 Pythagorean Triples





# Appendix D


## Boring stuff

### D.1 Version History

- v **0.1 2016:** This project is begun in a trio of physical exercise books as *The Little Book of Physics Formulae*, *The Little Book of Mathematics Formulae*, and *The Little Book of Astronomy Formulae*
- v **0.6 2016:** The process of transferring the formulae from paper to Latex is initiated, but abandoned (or drifted away from).
- v **0.7 2018-03-20:** The project is resurrected (probably because the author started MRes), uploaded to Overleaf, and cleaned up.
- v **0.8 2018-07-12:** Remaining formulae imported from the original books.
- v **0.9 2018-07-26:** Formulae imported from undergrad formula sheets.
- v **1.0 2018-07-28:** First public release, with some additions from 0.9.
- v **1.0.1 2018-07-31:** Minor corrections, added "dynamical timescale" (2.2.2) and some more formulae to the Statistics chapter (it was looking a little bare)

### D.2 Licensing

This work is licensed under a [Creative Commons "Attribution-ShareAlike 4.0 International"](#) license.



(Basically, as long as you credit me and share under a similar license, feel free to use this however you want)

### D.3 Contact

Visit [www.webofworlds.net](http://www.webofworlds.net) for science fiction, science fact, geeky opinions, and maybe some Python code.

Suggestions or corrections are welcome at [webofworlds@gmail.com](mailto:webofworlds@gmail.com)

### D.4 Credits

**Unit Circle:** By Jim.belk [CC BY-SA 3.0 (<https://creativecommons.org/licenses/by-sa/3.0>) or GFDL (<http://www.gnu.org/copyleft/fdl.html>)], from Wikimedia Commons ([https://commons.wikimedia.org/wiki/File:Unit\\_circle\\_angles\\_color.svg](https://commons.wikimedia.org/wiki/File:Unit_circle_angles_color.svg))

**Periodic Table:** By Dmarcus100 [CC BY-SA 4.0 (<https://creativecommons.org/licenses/by-sa/4.0>)], from Wikimedia Commons ([https://commons.wikimedia.org/wiki/File:Periodic\\_Table\\_Of\\_Elements\\_Atomic\\_Mass\\_Black\\_And\\_White.jpg](https://commons.wikimedia.org/wiki/File:Periodic_Table_Of_Elements_Atomic_Mass_Black_And_White.jpg))

1 <b>H</b> Hydrogen 1.008	4 <b>Be</b> Beryllium 9.012	10 <b>Ne</b> Neon 20.180	2 <b>He</b> Helium 4.003
3 <b>Li</b> Lithium 6.94	12 <b>Mg</b> Magnesium 24.305	18 <b>Ar</b> Argon 39.948	10 <b>Ne</b> Neon 20.180
11 <b>Na</b> Sodium 22.990	20 <b>Ca</b> Calcium 40.078	17 <b>Cl</b> Chlorine 35.45	9 <b>F</b> Fluorine 18.998
19 <b>K</b> Potassium 39.098	21 <b>Sc</b> Scandium 44.956	35 <b>Br</b> Bromine 79.904	36 <b>Kr</b> Krypton 83.798
37 <b>Rb</b> Rubidium 85.468	38 <b>Sr</b> Strontium 87.62	53 <b>I</b> Iodine 126.904	54 <b>Xe</b> Xenon 131.293
55 <b>Cs</b> Cesium 132.905	56 <b>Ba</b> Barium 137.327	83 <b>Bi</b> Bismuth 208.980	86 <b>Rn</b> Radon [222]
87 <b>Fr</b> Francium [223]	88 <b>Ra</b> Radium [226]	115 <b>Mc</b> Moscovium [289]	118 <b>Og</b> Oganesson [294]
	* 57 - 70	71 <b>Lu</b> Lutetium 174.967	71 <b>Lu</b> Lutetium 174.967
	** 89 - 102	103 <b>Lr</b> Lawrencium [262]	103 <b>Lr</b> Lawrencium [262]
		72 <b>Hf</b> Hafnium 178.49	72 <b>Hf</b> Hafnium 178.49
		73 <b>Ta</b> Tantalum 180.948	73 <b>Ta</b> Tantalum 180.948
		74 <b>W</b> Tungsten 183.84	74 <b>W</b> Tungsten 183.84
		75 <b>Re</b> Rhenium 186.207	75 <b>Re</b> Rhenium 186.207
		76 <b>Os</b> Osmium 190.23	76 <b>Os</b> Osmium 190.23
		78 <b>Ir</b> Iridium 192.217	78 <b>Ir</b> Iridium 192.217
		79 <b>Pt</b> Platinum 195.084	79 <b>Pt</b> Platinum 195.084
		80 <b>Au</b> Gold 196.967	80 <b>Au</b> Gold 196.967
		81 <b>Hg</b> Mercury 200.592	81 <b>Hg</b> Mercury 200.592
		82 <b>Pb</b> Lead 207.2	82 <b>Pb</b> Lead 207.2
		83 <b>Bi</b> Bismuth 208.980	83 <b>Bi</b> Bismuth 208.980
		84 <b>Po</b> Polonium [209]	84 <b>Po</b> Polonium [209]
		85 <b>At</b> Astatine [210]	85 <b>At</b> Astatine [210]
		116 <b>Lv</b> Livermorium [293]	116 <b>Lv</b> Livermorium [293]
		117 <b>Ts</b> Tennessine [293]	117 <b>Ts</b> Tennessine [293]
		118 <b>Og</b> Oganesson [294]	118 <b>Og</b> Oganesson [294]
		21 <b>Sc</b> Scandium 44.956	21 <b>Sc</b> Scandium 44.956
		22 <b>Ti</b> Titanium 47.867	22 <b>Ti</b> Titanium 47.867
		23 <b>V</b> Vanadium 50.942	23 <b>V</b> Vanadium 50.942
		24 <b>Cr</b> Chromium 51.996	24 <b>Cr</b> Chromium 51.996
		25 <b>Mn</b> Manganese 54.938	25 <b>Mn</b> Manganese 54.938
		26 <b>Fe</b> Iron 55.845	26 <b>Fe</b> Iron 55.845
		27 <b>Co</b> Cobalt 58.933	27 <b>Co</b> Cobalt 58.933
		28 <b>Ni</b> Nickel 58.693	28 <b>Ni</b> Nickel 58.693
		29 <b>Cu</b> Copper 63.546	29 <b>Cu</b> Copper 63.546
		30 <b>Zn</b> Zinc 65.38	30 <b>Zn</b> Zinc 65.38
		31 <b>Ga</b> Gallium 69.723	31 <b>Ga</b> Gallium 69.723
		32 <b>Ge</b> Germanium 72.630	32 <b>Ge</b> Germanium 72.630
		33 <b>As</b> Arsenic 74.922	33 <b>As</b> Arsenic 74.922
		34 <b>Se</b> Selenium 78.97	34 <b>Se</b> Selenium 78.97
		35 <b>Br</b> Bromine 79.904	35 <b>Br</b> Bromine 79.904
		49 <b>In</b> Indium 114.818	49 <b>In</b> Indium 114.818
		50 <b>Sn</b> Tin 118.710	50 <b>Sn</b> Tin 118.710
		51 <b>Sb</b> Antimony 121.760	51 <b>Sb</b> Antimony 121.760
		53 <b>Te</b> Tellurium 127.60	53 <b>Te</b> Tellurium 127.60
		81 <b>Tl</b> Thallium 204.38	81 <b>Tl</b> Thallium 204.38
		82 <b>Pb</b> Lead 207.2	82 <b>Pb</b> Lead 207.2
		83 <b>Bi</b> Bismuth 208.980	83 <b>Bi</b> Bismuth 208.980
		113 <b>Nh</b> Nihonium [286]	113 <b>Nh</b> Nihonium [286]
		114 <b>Fl</b> Flerovium [289]	114 <b>Fl</b> Flerovium [289]
		115 <b>Mc</b> Moscovium [289]	115 <b>Mc</b> Moscovium [289]
		116 <b>Lv</b> Livermorium [293]	116 <b>Lv</b> Livermorium [293]
		117 <b>Ts</b> Tennessine [293]	117 <b>Ts</b> Tennessine [293]
		118 <b>Og</b> Oganesson [294]	118 <b>Og</b> Oganesson [294]
		41 <b>Nb</b> Niobium 92.906	41 <b>Nb</b> Niobium 92.906
		42 <b>Mo</b> Molybdenum 95.95	42 <b>Mo</b> Molybdenum 95.95
		43 <b>Tc</b> Technetium [97]	43 <b>Tc</b> Technetium [97]
		44 <b>Ru</b> Ruthenium 101.07	44 <b>Ru</b> Ruthenium 101.07
		45 <b>Rh</b> Rhodium 102.906	45 <b>Rh</b> Rhodium 102.906
		46 <b>Pd</b> Palladium 106.42	46 <b>Pd</b> Palladium 106.42
		47 <b>Ag</b> Silver 107.868	47 <b>Ag</b> Silver 107.868
		48 <b>Cd</b> Cadmium 112.414	48 <b>Cd</b> Cadmium 112.414
		77 <b>Ir</b> Iridium 192.217	77 <b>Ir</b> Iridium 192.217
		78 <b>Pt</b> Platinum 195.084	78 <b>Pt</b> Platinum 195.084
		79 <b>Au</b> Gold 196.967	79 <b>Au</b> Gold 196.967
		80 <b>Hg</b> Mercury 200.592	80 <b>Hg</b> Mercury 200.592
		81 <b>Tl</b> Thallium 204.38	81 <b>Tl</b> Thallium 204.38
		82 <b>Pb</b> Lead 207.2	82 <b>Pb</b> Lead 207.2
		83 <b>Bi</b> Bismuth 208.980	83 <b>Bi</b> Bismuth 208.980
		105 <b>Db</b> Dubnium [270]	105 <b>Db</b> Dubnium [270]
		106 <b>Sg</b> Seaborgium [269]	106 <b>Sg</b> Seaborgium [269]
		107 <b>Bh</b> Bohrium [270]	107 <b>Bh</b> Bohrium [270]
		108 <b>Hs</b> Hassium [270]	108 <b>Hs</b> Hassium [270]
		109 <b>Mt</b> Meitnerium [278]	109 <b>Mt</b> Meitnerium [278]
		110 <b>Ds</b> Darmstadtium [281]	110 <b>Ds</b> Darmstadtium [281]
		111 <b>Rg</b> Roentgenium [281]	111 <b>Rg</b> Roentgenium [281]
		112 <b>Cn</b> Copernicium [285]	112 <b>Cn</b> Copernicium [285]
		113 <b>Nh</b> Nihonium [286]	113 <b>Nh</b> Nihonium [286]
		114 <b>Fl</b> Flerovium [289]	114 <b>Fl</b> Flerovium [289]
		115 <b>Mc</b> Moscovium [289]	115 <b>Mc</b> Moscovium [289]
		116 <b>Lv</b> Livermorium [293]	116 <b>Lv</b> Livermorium [293]
		117 <b>Ts</b> Tennessine [293]	117 <b>Ts</b> Tennessine [293]
		118 <b>Og</b> Oganesson [294]	118 <b>Og</b> Oganesson [294]
		39 <b>Y</b> Yttrium 88.906	39 <b>Y</b> Yttrium 88.906
		40 <b>Zr</b> Zirconium 91.224	40 <b>Zr</b> Zirconium 91.224
		41 <b>Nb</b> Niobium 92.906	41 <b>Nb</b> Niobium 92.906
		42 <b>Mo</b> Molybdenum 95.95	42 <b>Mo</b> Molybdenum 95.95
		43 <b>Tc</b> Technetium [97]	43 <b>Tc</b> Technetium [97]
		44 <b>Ru</b> Ruthenium 101.07	44 <b>Ru</b> Ruthenium 101.07
		45 <b>Rh</b> Rhodium 102.906	45 <b>Rh</b> Rhodium 102.906
		46 <b>Pd</b> Palladium 106.42	46 <b>Pd</b> Palladium 106.42
		47 <b>Ag</b> Silver 107.868	47 <b>Ag</b> Silver 107.868
		48 <b>Cd</b> Cadmium 112.414	48 <b>Cd</b> Cadmium 112.414
		49 <b>In</b> Indium 114.818	49 <b>In</b> Indium 114.818
		50 <b>Sn</b> Tin 118.710	50 <b>Sn</b> Tin 118.710
		51 <b>Sb</b> Antimony 121.760	51 <b>Sb</b> Antimony 121.760
		52 <b>Te</b> Tellurium 127.60	52 <b>Te</b> Tellurium 127.60
		53 <b>I</b> Iodine 126.904	53 <b>I</b> Iodine 126.904
		54 <b>Xe</b> Xenon 131.293	54 <b>Xe</b> Xenon 131.293
		71 <b>Lu</b> Lutetium 174.967	71 <b>Lu</b> Lutetium 174.967
		72 <b>Hf</b> Hafnium 178.49	72 <b>Hf</b> Hafnium 178.49
		73 <b>Ta</b> Tantalum 180.948	73 <b>Ta</b> Tantalum 180.948
		74 <b>W</b> Tungsten 183.84	74 <b>W</b> Tungsten 183.84
		75 <b>Re</b> Rhenium 186.207	75 <b>Re</b> Rhenium 186.207
		76 <b>Os</b> Osmium 190.23	76 <b>Os</b> Osmium 190.23
		77 <b>Ir</b> Iridium 192.217	77 <b>Ir</b> Iridium 192.217
		78 <b>Pt</b> Platinum 195.084	78 <b>Pt</b> Platinum 195.084
		79 <b>Au</b> Gold 196.967	79 <b>Au</b> Gold 196.967
		80 <b>Hg</b> Mercury 200.592	80 <b>Hg</b> Mercury 200.592
		81 <b>Tl</b> Thallium 204.38	81 <b>Tl</b> Thallium 204.38
		82 <b>Pb</b> Lead 207.2	82 <b>Pb</b> Lead 207.2
		83 <b>Bi</b> Bismuth 208.980	83 <b>Bi</b> Bismuth 208.980
		103 <b>Lr</b> Lawrencium [262]	103 <b>Lr</b> Lawrencium [262]
		104 <b>Rf</b> Rutherfordium [261]	104 <b>Rf</b> Rutherfordium [261]
		105 <b>Db</b> Dubnium [270]	105 <b>Db</b> Dubnium [270]
		106 <b>Sg</b> Seaborgium [269]	106 <b>Sg</b> Seaborgium [269]
		107 <b>Bh</b> Bohrium [270]	107 <b>Bh</b> Bohrium [270]
		108 <b>Hs</b> Hassium [270]	108 <b>Hs</b> Hassium [270]
		109 <b>Mt</b> Meitnerium [278]	109 <b>Mt</b> Meitnerium [278]
		110 <b>Ds</b> Darmstadtium [281]	110 <b>Ds</b> Darmstadtium [281]
		111 <b>Rg</b> Roentgenium [281]	111 <b>Rg</b> Roentgenium [281]
		112 <b>Cn</b> Copernicium [285]	112 <b>Cn</b> Copernicium [285]
		113 <b>Nh</b> Nihonium [286]	113 <b>Nh</b> Nihonium [286]
		114 <b>Fl</b> Flerovium [289]	114 <b>Fl</b> Flerovium [289]
		115 <b>Mc</b> Moscovium [289]	115 <b>Mc</b> Moscovium [289]
		116 <b>Lv</b> Livermorium [293]	116 <b>Lv</b> Livermorium [293]
		117 <b>Ts</b> Tennessine [293]	117 <b>Ts</b> Tennessine [293]
		118 <b>Og</b> Oganesson [294]	118 <b>Og</b> Oganesson [294]
		57 <b>La</b> Lanthanum 138.905	57 <b>La</b> Lanthanum 138.905
		58 <b>Ce</b> Cerium 140.116	58 <b>Ce</b> Cerium 140.116
		59 <b>Pr</b> Praseodymium 140.908	59 <b>Pr</b> Praseodymium 140.908
		60 <b>Nd</b> Neodymium 144.242	60 <b>Nd</b> Neodymium 144.242
		61 <b>Pm</b> Promethium [145]	61 <b>Pm</b> Promethium [145]
		62 <b>Sm</b> Samarium 150.36	62 <b>Sm</b> Samarium 150.36
		63 <b>Eu</b> Europium 151.964	63 <b>Eu</b> Europium 151.964
		64 <b>Gd</b> Gadolinium 157.25	64 <b>Gd</b> Gadolinium 157.25
		65 <b>Tb</b> Terbium 158.925	65 <b>Tb</b> Terbium 158.925
		66 <b>Dy</b> Dysprosium 162.500	66 <b>Dy</b> Dysprosium 162.500
		67 <b>Ho</b> Holmium 164.930	67 <b>Ho</b> Holmium 164.930
		68 <b>Er</b> Erbium 167.259	68 <b>Er</b> Erbium 167.259
		69 <b>Tm</b> Thulium 168.934	69 <b>Tm</b> Thulium 168.934
		70 <b>Yb</b> Ytterbium 173.045	70 <b>Yb</b> Ytterbium 173.045
		89 <b>Ac</b> Actinium [227]	89 <b>Ac</b> Actinium [227]
		90 <b>Th</b> Thorium 232.038	90 <b>Th</b> Thorium 232.038
		91 <b>Pa</b> Protactinium 231.036	91 <b>Pa</b> Protactinium 231.036
		92 <b>U</b> Uranium 238.029	92 <b>U</b> Uranium 238.029
		93 <b>Np</b> Neptunium [237]	93 <b>Np</b> Neptunium [237]
		94 <b>Pu</b> Plutonium [244]	94 <b>Pu</b> Plutonium [244]
		95 <b>Am</b> Americium [243]	95 <b>Am</b> Americium [243]
		96 <b>Cm</b> Curium [247]	96 <b>Cm</b> Curium [247]
		97 <b>Bk</b> Berkelium [247]	97 <b>Bk</b> Berkelium [247]
		98 <b>Cf</b> Californium [251]	98 <b>Cf</b> Californium [251]
		99 <b>Es</b> Einsteinium [252]	99 <b>Es</b> Einsteinium [252]
		100 <b>Fm</b> Fermium [257]	100 <b>Fm</b> Fermium [257]
		101 <b>Md</b> Mendelevium [258]	101 <b>Md</b> Mendelevium [258]
		102 <b>No</b> Nobelium [259]	102 <b>No</b> Nobelium [259]

\*Lanthanide series

\*\*Actinide series